



Determinants of Medicinal Plants Usage for Traditional Herbal Medicine among Villagers in Ibadan, Oyo State, Nigeria

O. C. Ariyo^{1*} and M. O. Ariyo²

¹Department of Vocational and Technical Studies, Forestry Research Institute of Nigeria, Federal College of Forestry Mechanization, Afaka, Kaduna, Kaduna State, Nigeria.

²Department of Horticulture and Landscape Technology, Forestry Research Institute of Nigeria, Federal College of Forestry Mechanization, Afaka, Kaduna, Kaduna State, Nigeria.

Authors' contributions

This work was carried out in collaboration between both authors. Author OCA designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. Authors OCA and MOA managed the analyses of the study. Author MOA managed the literature searches. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJRAF/2018/40141

Editor(s):

(1) Md. Abiar Rahman, Department of Agroforestry and Environment, Bangabandhu Sheikh Mujibur Rahman- Agricultural University (BSMRAU), Bangladesh.

Reviewers:

- (1) Esraa Ashraf Ahmed ElHawary, Ain Shams University, Egypt.
(2) Muhammad Ali, Kano University of Science and Technology, Nigeria.
(3) Izharul Hasan, India.

Complete Peer review History: <http://www.sciencedomain.org/review-history/23859>

Original Research Article

Received 13th January 2018
Accepted 20th March 2018
Published 28th March 2018

ABSTRACT

The study was designed to access the determinants of medicinal plants for traditional herbal medicine among villagers living at the perimeter fence of International Institute of Tropical Agriculture (IITA), Ibadan, Oyo State, Nigeria. The multistage sampling procedure was adopted for the study. A total number of four hundred and eight respondents comprising of farmers, hunters, herbalists and herb sellers were randomly selected and interviewed using copies of a well-structured questionnaire. Data were analysed using descriptive statistics, Probit and Tobit regression analysis. The study showed that the average ages of farmers and hunters were 55 and 57 years while the average age of herb sellers was 43 and herbalist 63 years. Majority of the respondents pooled together were males, married with an average age of 55 years and household

*Corresponding author: E-mail: ask4ariyo@yahoo.com;

size of 7 members. The larger percentage of them were native of the study area, not educated, not employed, but having the monthly income between 12,000- 20,000 naira and closer to the forest by 1-9 km. The study further revealed that there was a significant relationship between the use and intensity of use of medicinal plants for traditional herbal medicine and factors that determine it. Variables such as age, religion, sex, believe in traditional herbs, forest medicinal plants used in treating any ailment in the past, nearness to the forest, the presence of healthcare medical centre, poverty status and income were significant at 1% probability level. Household size and occupation was significant at 5% level while the location was significant at 10% level of significance. The study, therefore, recommends that conservation and domestication of these valuable medicinal plants should be a priority to prevent their extinction and ensure their continues supply to people that need them.

Keywords: Medicinal plants; traditional herbal medicine; uses, variables; villages; respondents; stratified; probit and tobit.

1. INTRODUCTION

Human beings, through intuition, have invented the art of healing systems, which are mostly based on medicinal plants. These medicinal plants are accredited with mystical and supernatural powers of healing. They are used widely across the world for primary health care and also in modern drug discoveries. It is estimated that more than 13,000 species of Medicinal and Aromatic Plants (MAPs) are used in traditional medicines and herbal cosmetics throughout the world [1].

Herbal medicine has been defined differently by various people. According to [2], "herbal drugs constitute only those traditional medicines which primarily use medicinal plant preparations for their therapy". [3] also defines herbal medicine as "the use of plant products to treat or prevent disease". [4] suggests that the treatment of herbal practitioners usually "takes the form of herbs, plant preparations, and prayers. The World Health Organization (WHO) defines herbal medicine as "a plant-derived material or preparation with therapeutic or other human health benefits which contains either raw or processed ingredients from one or more plants ([5], [6]). However, the WHO Regional Office for Africa (2004) uses the term "traditional medicine" as a synonym for herbal medicine and defines it as "the use of indigenous medicinal and aromatic plants, animal parts, or organic and inorganic materials for preventive and therapeutic purposes.

Traditional medicine continues to play an essential role in improving and maintaining health in developing countries [7]. Traditional medicine is defined as 'diverse health practices, approaches, knowledge and beliefs incorporating

plant, animal, and/ or mineral based medicine, spiritual therapies, manual techniques and exercises applied singularly or in combination to maintain well-being, as well as to treat, diagnose or prevent illness' [8,9,10]. Policies for the integration of traditional medicine into public health care systems have to vary extents been formulated in some countries [11], and the share of the population using traditional medicine are reported to be as high as 40% in China and 80% in Africa [9]. In Nigeria, there has been a consistent and noticeable shift from the earlier preference in favour of orthodox medicine to greater acceptance of traditional (herbal) medicines as in many other countries worldwide [12]. Over 90% of Nigerians in rural areas and 40% in urban areas depend partly or wholly on traditional medicine [13].

Traditional medicine is generally considered highly available and accessible to people in developing countries [14]. This high use of herbal medicines may be due to accessibility, affordability, availability and acceptability of traditional medicines by a majority of the populace in developing countries [15]. Consequently, poor and marginalised people are commonly assumed to be most reliant on traditional medicine for their healthcare [7]. Differences in reliance have also been observed between rural and urban areas [16].

Traditional and herbal medicine has taken the new name, complementary and alternative medicine (CAM). CAM refers to those therapeutic and diagnostic disciplines that exist largely outside the institutions where orthodox or modern health care is provided [17]. On the other hand, the [18] defined herbal medicine as "plant's seeds, berries, roots, leaves, bark, or flowers for medicinal purposes. Herbs that are used for

medicinal purposes come in a variety of forms. Active parts of a plant may include leaves, flowers, stems, roots, seeds, and berries [19]. They may be taken internally as pills or powders, dissolved into tinctures or syrups, or brewed in teas and concoctions.

A medicinal plant is defined as any substance with one or more of its organ containing substances that can be used for therapeutic purposes or which can be used as precursors for the synthesis of antimicrobial drugs [20,21]. It is estimated that there are about 250, 000 – 500, 000 species of plants on earth [22], of which a relatively small percentage (1-10%) of these are used for food by humans and animals. It is possible that more serve medicinal purposes [23].

Medicinal plants contain numerous biologically active compounds such as carbohydrates, proteins, enzymes, fats and oils, minerals, vitamins, alkaloids, quinones, terpenoids, flavonoids, carotenoids, sterols, simple phenolic glycosides, tannins, saponins, polyphenols, to mention a few which have medicinal activities.

In African countries, approximately 80% of the population uses traditional medicine for the treatment of various diseases and ailments like malaria, typhoid, ulcer, skin diseases, diabetes, reproductive problems, aches and pains for various socio-cultural & economic reasons [24]. In Nigeria, the majority of citizens still uses medicinal plants and visit traditional medicine practitioners for their health care need [25]. It was reported by WHO that in Nigeria, the ratio of Traditional Health Practitioners to the population was 1:110, while the ratio of Medical Doctors to the population was 1:16, 400 [26]. This gives credence to the fact that people patronise. Traditional medicine practitioners (TMPs) for their primary health needs more than orthodox medical doctors.

In Nigeria, the use of medicinal plants for traditional and/or herbal remedies have become more popular in the treatment of minor ailments, and also on account of the increasing costs of prescription drugs in the maintenance of personal health and well-being, and the bioprospecting of new plant-derived drugs. Based on current research and financial investments, medicinal plants will, seemingly, continue to play an essential role as a health aid ([27]; [23]). In spite of the millions of chemical compounds currently synthesised in the laboratory, and available for screening for an

action of therapeutic value, natural products, particularly of plants origin remain the most important sources of new drugs [25]. Indeed, the market and public demand for these traditional and herbal remedies has been so great that there is a significant risk that many medicinal plants today, face either extinction or loss of genetic diversity. The factors that determine the use of medicinal plants for traditional herbal medicine are not known, it is on this premise that this research was carried out to access the determinants of medicinal plants for traditional herbal medicine among respondents of the study area.

Specific objectives are to:

- i. describe the socio-economic characteristics of the respondents.
- ii. determine the factors that affect the use of medicinal plants for traditional herbal medicine by the respondents and
- iii. determine the level or intensity of medicinal plants usage for traditional herbal medicine by the respondents.

1.1 The Hypothesis of the Study

The hypothesis of the study was stated in the null form as follows:

Ho: There was no significant relationship between the medicinal plants usage for traditional herbal medicine and the factors that determine it.

Ho: There was no significant relationship between the intensity of medicinal plant usage for traditional herbal medicine and the factors that determine it.

2. METHODOLOGY

2.1 Study Area

The study areas are the villages by IITA perimeter fence in Akinyele Local Government area of Ibadan, Oyo State, Nigeria. IITA is located at longitude 7° 30' 8"N, latitude 3° 54' 37"E and 243 m above sea level. In 1965, the Federal Government of Nigeria allocated some 1000 hectares of land for the establishment of the main IITA campus. Prior to the acquisition of land by IITA through the Federal Government of Nigeria, there were patches of secondary forest which serves as a means of livelihood to the villagers in the area. The most extensive land

use pattern was arable and tree crop and about 3000 people lived in about twenty-eight villages scattered in this area. These villages were relocated to the perimeter fence of IITA where there was an expanse of secondary forest. At the period of this study, only seventeen villages exist at the perimeter fence of IITA and the secondary forest had been taken over by development leaving patches of scattered forest in the area.

2.2 Data Collection and Sampling Methods

A multistage sampling procedure was adopted for this study. All the seventeen villages by IITA perimeter fence were purposefully selected because of the following reasons (i) the villages were once located on the area where IITA is presently located (ii) the closeness of the villages to IITA forest and (iii) the presence of forest patches in all the villages. Respondents were stratified into four major groups in each of the village: namely farmers, hunters, herb sellers and herbalist. Within each stratum, a random selection of six respondents was carried out making twenty-four respondents in each village and a total number of four hundred and eight respondents in all the seventeen villages. An interview was conducted for each of the respondents with the aid of questionnaires and responses were recorded.

2.3 Data Analysis

Data were analysed using descriptive statistics such as tables, frequency, percentages and means to summarise the socioeconomics data. Probit model was used to determine factors affecting the use of medicinal plants for traditional or herbal medicine by the respondents and Tobit regression analysis was used following ([28]; [29]; [30]), whose works were built on [31] to determine the intensity of medicinal plant usage for traditional herbal medicine. Probit is an estimating model that emerges from the normal distribution function. It is useful in a regression that involves binary response of 0 and 1 [32].

The model is specified implicitly as follows:

$$P_1 = P_r (Y_1 = 1) = P_r (U_1 > U_{01}) = F_1 (X_1 \beta) \quad (1)$$

P_1 = Probit notation

X = Matrix of the explanatory variables included in the model

X_1 = Age of the respondents (Years),

X_2 = Believe in traditional herb (BTH) (Yes= 1, 0 otherwise),

X_3 = Religion (Traditional religion= 1, 0 otherwise),

X_4 = Sex (Male= 1; 0 otherwise),

X_5 = Forest medicinal plants used in treating any ailment in the past (FPUTAP) (Yes= 1, 0 otherwise),

X_6 = Nearness of respondents to the forest medicinal plants (Distance in km),

X_7 = Household size (Actual number of household members),

X_8 = Occupation

X_9 = Level of education (years of schooling),

X_{10} = Marital status

X_{11} = Location of the respondent (Rural area= 1, 0 otherwise),

X_{12} = Presence of health care centres (PHCC) - Hospital (Yes= 1, 0 otherwise),

X_{13} = Nativity of the household (native= 1, 0 otherwise),

X_{14} = Poverty status of the respondent (PSR) (Poor= 1, 0 otherwise).

X_{15} = Income (₦)

β = Vector of parameter to be estimated

P_r = Probability function of using medicinal plants for traditional herbal medicine (1, 0 otherwise)

$F (X_1 \beta)$ = Cumulative distribution function for random error term (U_1) evaluated at $X_1 \beta$

- Explicitly, the probit is specified as:

$$P_1 = P (FAD = 1/x) = b_0 + b_1 x_1 + b_2 x_2 + \dots + b_{15} x_{15} + e \quad (2)$$

Where $P (FAD = 1/x)$ = Probability decision of using medicinal plants for traditional or herbal medicine (1, or 0 otherwise)

$b_1 - b_{15}$ = Maximum likelihood estimates.

$X_1 - X_{15}$ = Explanatory variables as defined in the implicit form of the Probit model above.

e = error term

The conceptual Tobit model can be specified as;

$$y_i^* = X_i \beta + \varepsilon_i \quad (3)$$

$$y_i = y_i^* \text{ if } y_i^* > 0$$

$$y_i = 0 \text{ if } y_i^* \leq 0$$

y_i is the observed dependent variables indicating the level or intensity of use of medicinal plants, y_i^* is the latent dependent variables, x_i is the vector of the independent variable, β is the vector of coefficients, ε_i is assumed to be independently normally distributed: $\varepsilon \sim N(0, \delta)$ and therefore $y_i \sim N(X_i \beta, \delta)$.

The extent or intensity of utilization of medicinal plant (y_i) was measured using likert scale of Never = 0, rarely = 0.2, sometimes = 0.4, frequent = 0.6 and always = 0.8.

The explanatory (x_i) variables include:

- X_1 = Age of the respondents (Years),
- X_2 = Believe in traditional herb (BTH) (Yes= 1, 0 otherwise),
- X_3 = Religion (Traditional religion= 1, 0 otherwise),
- X_4 = Sex (Male= 1; 0 otherwise),
- X_5 = Nearness of respondents to the forest medicinal plants (Distance in km),
- X_6 = Distance of health services centers (DHSC) (Km),
- X_7 = Household size (Actual number of household members),
- X_8 = Occupation
- X_9 = Level of education (years of schooling),
- X_{10} = Marital status
- X_{11} = Location of the respondent (Rural area= 1, 0 otherwise),
- X_{12} = Nativity of the household (native= 1, 0 otherwise),
- X_{13} = Poverty status of the respondents (PSR) (Poor= 1, 0 otherwise).
- X_{14} = Income (₦)

3. RESULTS AND DISCUSSION

3.1 Socio-economic Characteristics of the Respondents

Table 1 showed the socio-economic characteristics of the respondents. The average age of farmers and hunters was 55 and 57 years while the average age of herb sellers was 43 and herbalist 63 years. The highest age group was found between 41- 60 years for farmers, hunters and herb sellers with 67.65%, 68.63% and 60.78% respectively while 64.71% of herbalist had the highest age between 61-80 years. The age of an individual determines his/ her utilization of medicinal plants for traditional herbal medicine. The use of medicinal plants increases with age.

The percentage of farmers that were male was 85.29 while 14.71% were female. 100% of hunters and herbalist were male while herb sellers had 100% female. 71.32% of the total respondents were male while 28.68% were female. A higher percentage of the males will use medicinal plants for traditional herbal medicine. This can be explained by a larger household size in male headed households, where medicinal

plants would have to be used for traditional herbal medicine as curative measure in order to minimize cost.

Majority of the respondents were married with hunters' respondents having the highest value of 96.08% followed by herbalists 95.10%, farmers and herb sellers had 94.12% and 92.16% respectively. 94.36% of the total respondents were married, 2.21% were single, 2.45% and 0.98% were widower and widowed respectively. The married respondents will use medicinal plants for traditional herbal medicine. This is explained by a large sized household in the case of married respondents where medicinal plants is preferred to reduce the amount of income spent on curative measures.

The highest household size was found in the group of 6-10 for all the categories of respondents, herbalist had the highest household size of 77.45%, followed by hunters, herb sellers and farmers with 71.57%, 64.71% and 60.78% respectively. 68.63% of the total respondents had household size between 6-10, 12.99% had family size within 11-15 while only 18.38% had it between 1-5. The larger the household size the greater the use of medicinal plants for traditional herbal medicine. A large household will prefers medicinal plants in treatment of ailments because of its affordability and the inability of the household to purchase prescriptive drugs.

The percentage of the respondents that were not educated was 97.06%, 86.27%, 68.63% and 67.65% for hunters, herbalist, herb sellers and famers. Only 22.55%, 21.57%, 12.75% and 1.96% farmers, herb sellers, herbalist and hunters had primary six educations while 9.80% of famers and herb sellers, and 0.98% of hunters and herbalist had secondary school education. The total number of respondents that were educated both primary and secondary school education was 20.10% while 79.90% of them were not educated. An individual with no or low level of education will use medicinal plants for traditional herbal medicine, this is due to the number of years for which the respondents has been exposed to formal education while an individual with higher level of education prefers orthodox medicine because of its ease of consumption, storability and carriage.

Majority of the respondents interviewed were native of the area with a value of 89.71% while 10.29% were non native residing in the area. The

Table 1. Socio- economic characteristics of respondents

Socio economics characteristics	Crop farmers		Hunters		Herb sellers		Herbalist		Total	Percentage
	Frequency	%age	Frequency	%age	Frequency	%age	Frequency	%age		
Age										
21-40	5	4.90	1	0.98	40	39.22	-	-	46	11.27
41-60	69	67.65	70	68.63	62	60.78	33	32.35	234	57.35
61-80	28	27.45	29	28.43	-	-	66	64.71	123	30.15
81-100	-	-	2	1.96	-	-	3	2.94	5	1.23
Sex										
Male	87	85.29	102	100	-	-	102	100	291	71.32
Female	15	14.71	-	-	102	100	-	-	117	28.68
Marital Status										
Single	-	-	1	0.98	8	7.84	-	-	9	2.21
Married	96	94.12	98	96.08	94	92.16	97	95.10	385	94.36
Widowed	4	3.92	-	-	-	-	-	-	4	0.98
Widower	2	1.96	3	2.94	-	-	5	4.90	10	2.45
Household size										
1-5	39	38.24	4	3.92	29	28.43	3	2.94	75	18.38
6-10	62	60.78	73	71.57	66	64.71	79	77.45	280	68.63
11-15	1	0.98	25	24.51	7	6.86	20	19.61	53	12.99
Level of Education										
Primary six	23	22.55	2	1.96	22	21.57	13	12.75	60	14.71
Secondary	10	9.80	1	0.98	10	9.80	1	0.98	22	5.39
Not educated	69	67.65	99	97.06	70	68.63	88	86.27	326	79.90
Nativity										
Native	92	90.20	87	85.29	98	96.08	89	87.25	366	89.71
Non-native	10	9.80	15	14.71	4	3.92	13	12.75	42	10.29
Nearness to forest (Km)										
1-3	88	86.27	38	37.25	68	66.67	85	83.33	279	68.38
4-6	14	13.73	49	48.04	25	24.51	12	11.77	100	24.50
7-9	-	-	15	14.71	9	8.82	5	4.90	29	7.12

Socio economics characteristics	Crop farmers		Hunters		Herb sellers		Herbalist		Total	Percentage
	Frequency	%age	Frequency	%age	Frequency	%age	Frequency	%age		
Occupation										
Farming	80	78.43	-	-	-	-	-	-	80	19.61
Hunting	-	-	88	86.27	-	-	-	-	88	21.57
Herb selling	-	-	-	-	77	75.49	-	-	77	18.87
Herbalist	-	-	-	-	-	-	96	94.12	96	23.53
Others	22	21.57	14	13.73	25	24.51	6	5.88	67	16.42
Employment										
Employed	21	20.59	62	60.78	-	-	13	12.75	96	23.53
Not employed	81	79.41	40	39.22	102	100	89	87.25	312	76.47
Income										
4000- 12,000	58	56.86	37	36.27	45	44.12	6	5.88	146	35.78
12,000- 20,000	44	43.14	60	58.82	57	55.88	21	20.59	182	44.61
20,000-28,000	-	-	5	4.90	-	-	54	52.94	59	14.46
28,000-36000	-	-	-	-	-	-	21	20.59	21	5.15

Source: Computed from Field Survey Data, 2016

native are expected to use medicinal plants for traditional herbal medicine than non native. This is because the native are well familiar with medicinal plants in their environments with their curative values.

The nearness of the respondents to the forest showed that 86.27% of famers, 83.33% of herbalist, 66.67% herb sellers, and 37.25% of the hunters were closer to the forest with a distance of 1-3 km. The percentage of hunters, herb sellers, herbalists and farmers that were closer to the forest by 4-6 km were 48.04%, 24.51%, 11.77% and 13.73% respectively. Only 14.71%, 8.82%, and 4.90% of hunters, herb sellers and herbalist were closer to the forest by 7-9 km. 68.38% of the total respondents were closer to forest by 1-3 km while 24.50% and 7.12% of them had forest closer to them by 4-6 and 7-9 km respectively. The closer the forest to the respondents the greater the use of medicinal plants for traditional herbal medicine

In term of employment, all the herb sellers' respondents were not employed apart from selling of herbal plants, they formed 100%. The percentage of unemployed herbalist, farmers and hunters were 87.25%, 79.41% and 39.22% respectively while 60.78% of hunters, 20.59% of famers and 12.75% of herbalists were employed. The results pooled together showed that 76.47% of the respondents were not employed while only 23.53% were employed. The unemployed respondents are likely to use medicinal plants for traditional herbal medicine because of its affordability and availability.

56.86% of famers, 44.12% of herb sellers, 36.27% of hunters and 5.88% of herbalists had income ranges between 4, 000 to 12, 000 naira. The percentage of hunters, herb sellers, famers and herbalist that had their income ranges between 12, 000 to 20, 000 naira were 58.82%, 55.88%, 43.14% and 20.59% respectively. Only 52.94% and 20.59% of herbalist had their income ranges from 20, 000 to 28, 000, and 12, 000 to 20 000 and 28, 000 to 36, 000 naira while 4.90% of hunters had it between 20, 000 to 28, 000 naira. 44.61% and 35.78% of the total respondents had their income ranges between 12, 000- 20, 000 and 4, 000- 12, 000 while 14.46% and 5.15% had it between 20, 000-28, 000 and 28, 000-36, 000 respectively. The lower the employment and income, the greater the use of medicinal plants for traditional herbal

medicine. This is because of relative cheapness of medicinal plants compared to orthodox drugs.

3.2 Determinants of Medicinal Plants Usage for Traditional Herbal Medicine by the Respondents

Table 2 showed the result of factors that affect the use of medicinal plants for traditional herbal medicine by the respondents. From the table, eight out of fifteen variables had significant coefficients. These include religion (X_3), sex (X_4), forest plant used in treating any ailment in the past (X_5), nearness to the forest (X_6), occupation (X_8), presence of health care centers (X_{12}), poverty status of the respondents (X_{14}) and income (X_{15}). The significant and positive determinants of medicinal plants usage for traditional herbal medicine by the respondents include sex, medicinal plants used in treating any ailment in the past, nearness to the forest, occupation, poverty status and income. In other words, enhancing these variables enhances the likelihood of respondents to utilize medicinal plants for traditional and herbal medicine. According to [33], age, sex, and accessibility were significant and positive determinants of consumer preference for medicinal plants in Oyo metropolis Nigeria. Furthermore, significant and negative determinant of respondents utilization of medicinal plants for traditional herbal medicine include religion and presence of health care centre. However reducing these variables will enhances the respondent's likelihood of utilizing medicinal plants for traditional herbal medicine. Also, according to [33], primary education and household head education are significant and negative determinants of consumer preference for medicinal plants in Oyo metropolis, Nigeria. The log-likelihood ratio (LR) statistics of the entire model was -144.54722 and was significant at 1% level of significance, meaning that the overall model was significant and the null hypothesis was rejected. The coefficients of significant variables are explained thus:

Religion (X_3) was significant at 1% level of significance. It has a negative relationship with medicinal plant usage for traditional herbal medicine (-1.014154) which implies that a non-traditional religion respondents are likely not to use medicinal plants for traditional herbal medicine.

Sex (X_4) was significant at 1% level of significance. It has a positive relationship with medicinal plant usage (0.4111003). This means

that a male respondent or a household headed by male is likely to use medicinal plants from the forest for traditional herbal medicine since such a household is likely to be large. This is because of the relative cheapness and availability of the medicinal plants.

Forest medicinal plants used in treating any ailment in the past (X_5) was significant at 1% level of significant. It has a positive (0.649201) relationship with medicinal plants usage. This means that respondents that have used medicinal plants in the treatment of a particular illness in the past are likely to continued using it.

Nearness of the respondents to the forest (X_6) was significant at 1% level of significant. It has positive (0.2661395) relationship with medicinal plant usage. This means that the closer the respondents to the forest, the greater the likelihood of using medicinal plants for traditional herbal medicine.

Occupation (X_8) was significant at 10% level of significant. It has positive (0.1385835) relationship to the medicinal plants usage. This means that the closer the relationship of respondent's occupation to the forest, the more likelihood will be the respondents to use forest medicinal plants for traditional herbal medicine.

The coefficient of educational level (X_9) was negative (-0.0208478) but not significant meaning that the level of education of an individual reduces the use of medicinal plants for traditional herbal medicine. This agrees with the findings of [33] which stated that the percentage of respondents that prefer medicinal plants is inversely proportional to the educational status; this is due to the number of years for which the respondents has been exposed to formal education. An individual with higher level of education prefers orthodox medicine because of its ease of consumption, storability and carriage. [34] stated that the use of traditional medicine significantly decreased with education of household head in the rural hills site as did use of medicinal plants in the rural hills site and the peri-urban site. [35] argued that formal education in developing countries is the strongest form of exposure to Western paradigms, and that educated people therefore tend to opt for allopathic medicine. Many other studies have documented this same trend ([36]; [37]; [38]). The insignificant effect of education on the use of medicinal plants for traditional herbal medicine by the respondents could be explained by the

relatively strong cultural ties to medicinal plants for traditional herbal medicine ([34]; [39]) and also could be due to the fact that majority of the respondents in the study area were not educated.

The presence of health care medical centre (X_{12}) was significant at 1% level of significant. It has a negative relationship (-0.4787627) according to the a priori expectation with the medicinal plants usage. This means that the presence of health care medical centre reduces the use of medicinal plants for traditional herbal medicine by the respondents.

Poverty status of respondents (X_{14}) was significant at 1% level of significant. It has a positive relationship (0.5132115) according to the a priori expectation with the medicinal plants usage. This means that the poorer the respondents the more the use of medicinal plants for traditional herbal medicine. [34] stated that richer households were more likely to use traditional medicine and medicinal plants in the peri-urban site and traditional medicine in the rural mountain site. Studies in developed countries have reported increased use of alternative medicine with rising incomes [34]. For the rural hills site, [34] found a decreased likelihood of medicinal plant use for rich households, supporting the common assumption that traditional medicine is mostly relied on by the poor and disadvantaged ([35]; [40]; [9]).

The income of the respondents (X_{15}) was significant at 1% level of significant. It has a positive relationship as against the a priori expectation. This means that the higher the income the greater the use of medicinal plants for traditional herbal medicine by the respondents. According to [41], income is a strong predictor of herbal medicine utilization and is positively related to it. In most cases, having a lower income increases the use of herbal medicine, especially in developing countries like Ghana ([42]; [43]; [44]). On the hand, having a higher income increases the use of herbal medicine, especially in developed countries ([45]; [46]). This is because herbal drugs are relatively expensive in developed countries.

3.3 Determinants of the Level or Intensity of Medicinal Plants Usage for Traditional Herbal Medicine by the Respondents

This was analyzed by specifying and estimating a Tobit regression model. From table 3, nine

variables determined the intensity of medicinal plants usage by the respondents. The variables are age (X_1), believe in traditional herb (X_2), religion (X_3), sex (X_4), nearness to the forest (X_5), household size (X_7), occupation (X_8), location of the respondents (X_{11}), and poverty status of the respondents (X_{13}). According to [41], age, level of education and place of residence were background factors associated with the use of herbal medicine. The coefficient of variable X_1 , X_2 , X_5 , X_8 , X_{11} and X_{13} were positive and significant at 1% level while X_8 was significant at 5% and X_{11} was significant at 10% probability level. This was in accordance with a priori expectation that as the rate of these variables increases, the intensity of medicinal plants usage for traditional herbal medicine will also increase.

The coefficient of X_3 (religion) and X_4 (sex) were equally significant at 1% level while variables X_7 (household size) was significant at 5% probability level but were negatively signed meaning that as these variables increase, the intensity of medicinal plants usage for traditional herbal medicine decreases. The coefficients of significant variables are further explained thus:

Coefficient of age (X_1) was positive sign as expected and significant at 1 percent probability level. The higher the age the greater the experience and the more the tendency of

increasing the use of medicinal plants for traditional herbal medicine.

Coefficient of believe of the respondents in traditional herb represented by variable X_2 was positive sign as expected and significant at 1 percent probability level. Believe of the respondents in traditional herbs increases the intensity of use of medicinal plants.

Coefficient of religion (X_3) was negative sign as expected and significant at 1% probability level. The non-traditional religions are less expected to use medicinal plants for traditional herbal medicine.

Coefficient of sex (X_4) was negative sign and significant at 1% probability level. This agreed with the findings of [47] and [48] who found a significant relationship between sex and the use of herbal medicine.

The coefficient of nearness of the respondents to the forest (X_5) was positive sign as expected and significant at 1% probability level. The nearer the respondents to the forest the greater the likelihood of increasing the use of medicinal plants for traditional herbal medicine.

Coefficient of household size (X_7) was negative sign and significant at 5% probability level. This shows that an increase in household size decrease the intensity of use of forest medicinal

Table 2. Analysis of determinants of medicinal plants usage for traditional herbal medicine by the respondents using probit model

Variables	Coefficient	Standard error	Z values	P> Z values
Age (X_1)	-0.0071317	0.0107504	-0.66	0.507
BTH (X_2)	2.18e-06	0.0000424	0.05	0.959
Religion (X_3)	-1.014154	0.275508	-3.68	0.000***
Sex (X_4)	0.4111003	0.1117633	3.68	0.000***
FPUTAP (X_5)	0.649201	0.1763329	3.68	0.000***
Nearness to forest (X_6)	0.2661395	0.0724038	3.68	0.000***
House hold size (X_7)	0.0326353	0.0436195	0.75	0.454
Occupation (X_8)	0.1385835	0.0728342	1.90	0.057**
Level of education (X_9)	-0.0208478	0.0245616	-0.85	0.396
Marital status (X_{10})	0.1185416	0.322819	0.37	0.713
Location of respondents (X_{11})	-0.652795	0.4480325	-1.46	0.145
PHCC (X_{12})	-0.4787627	0.1302914	-3.67	0.000***
Nativity of the household (X_{13})	-0.0001033	0.0000965	-1.07	0.285
Poverty status of respondents (X_{14})	0.5132115	0.1395006	3.68	0.000***
Income (X_{15})	0.0001362	0.0000238	5.73	0.000***
Constant	-1.137793	0.9822856	-1.16	0.247

Source: Computed from Field Survey Data, 2016

*** Significant at 0.01, ** Significant at 0.05; Prob. >Chi2= 0.0000, LR chi2 (16) = 67.70, Pseudo R² = 0.1897, Log likelihood = -144.54722, Number of obs. = 408

Table 3. Analysis of determinants of the level or intensity of medicinal plants usage for traditional herbal medicine by the respondents using Tobit model

Variables	Coefficient	Standard error	T values	P> Z values
Age (X ₁)	0.003883	0.0009624	4.03	0.000***
BTH (X ₂)	0.2341244	0.0375628	6.23	0.000***
Religion (X ₃)	-0.2015844	0.0311281	-6.48	0.000***
Sex (X ₄)	-0.408688	0.0254503	-16.06	0.000***
Nearness to forest (X ₅)	0.1244676	0.0263149	4.73	0.000***
DHSC (X ₆)	-0.0014127	0.0037257	-0.38	0.705
House hold size (X ₇)	-0.0076084	0.0032877	-2.31	0.021**
Occupation (X ₈)	0.0153784	0.0067273	2.29	0.023**
Level of education (X ₉)	0.0030982	0.0020917	1.48	0.139
Marital status (X ₁₀)	0.0216682	0.0315034	0.69	0.492
Location of respondents (X ₁₁)	0.0494329	0.0296905	1.66	0.097*
Nativity of the household (X ₁₂)	-0.0094667	0.0295594	-0.32	0.749
Poverty status of respondents (X ₁₃)	0.1368722	0.0295149	4.64	0.000***
Income (X ₁₄)	2.19e-06	1.99e-06	1.10	0.272
Constant	0.4106552	0.0953051	4.31	0.000
Sigma	0.1300151	0.0046118		

Source: Computed from Field Survey Data, 2016.

*** Significant at 0.01, ** Significant at 0.05, * Significant at 0.10; Prob. >Chi2= 0.0000, LR chi2 (15) = 571.03, Pseudo R² = 7.1881, Log likelihood = 245.79393, Number of obs. = 408

plants. Household head with formal education is associated with lower level of preference for medicinal plant usage for traditional herbal medicine. [49] reports that households headed by females have limited access to medicinal plants when those are collected far from the house.

Coefficient of occupation (X₈) was positive sign as expected and significant at 5% probability level. The closer the relationship of respondent's occupation to the forest, the more likelihood will be the intensity of use of forest medicinal plants for traditional herbal medicine.

The coefficient of location of the respondents (X₁₁) was positive sign as expected and significant at 10% probability level; this conforms to the statement made by [50] and [43] that herbal medicine is often used in the rural areas than the urban areas. This is because rural residents have little or no access to orthodox medicine and find herbal medicine to be relatively less expensive ([50]; [43]). Traditional medicine use is more prevalent in rural than in town areas [16]. Also, [41] found a significant relationship between place of residence and the use of herbal medicine.

Coefficient of poverty status (X₁₃) was positive sign as expected and significant at 1 percent probability level. The poor respondents are expected to increase the use of forest medicinal

plants for traditional herbal medicine. This is also because the medicinal plants are relatively cheap and available.

4. CONCLUSION AND RECOMMENDATION

From the results of the study, it can be concluded that all the hunters' respondents are male and herb sellers are female while farmers and herbalist respondents are both male and female respectively. The average age of farmers and hunters was 55 and 57 years while the average age of herb sellers was 43 and herbalist 63 years. Majority of the respondents pooled together are male, married with average age of 55 years and house hold size of 7 members. The larger percentage of them were native of the study area, not educated, not employed, but having the monthly income between 12,000-20,000 naira and closer to the forest by 1-9 km.

The most important factors that affect the use of medicinal plants for traditional herbal medicine by the respondents are religion, sex, forest plant used in treating any ailment in the past, nearness to the forest, occupation, presence of health care centers, poverty status of the respondents and income while the level or intensity of medicinal plants usage for traditional herbal medicine were determined by factors such as age, believe in traditional herb, religion, sex, nearness to the forest, household size, occupation, location of

the respondents and poverty status of the respondents.

It is therefore recommended that conservation of these valuable medicinal plants should be a priority for the well-being and livelihoods of indigenous local communities and the society at large, which depend on these medicinal plants. Moreover, studies into the domestication of these medicinal plants, the types of plants, parts of plants used, efficacy and diseases cured should be investigated in the future. The documentation of plants and their therapeutic properties is an area that must be of interest to future researchers.

CONSENT

As per international standard or university standard, patient's written consent has been collected and preserved by the authors.

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the authors.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Wilkinson JA, Wahlqvist ML, Clark J. New food and pharmaceutical products from agriculture. rural industries research and development corporation, Australia. 2002:1-30.
2. Kamboj VP. Herbal medicine. *Curr. Scie.* 2000;78(1):35-51.
3. Lucas GN. Herbal medicine and children. *Sri Lanka Journal of Child Health.* 2010;39:76-78.
4. Nsawah-Nuamah NNN, Overbosch B, Boom van den GJM. Health care provision and self - Medication in Ghana. Legon: University of Ghana. ISSER; 2004. (Accessed from on 14 September 2010) Available:[http://www.isser.org/publications/older/healthcare provision](http://www.isser.org/publications/older/healthcare%20provision).
5. WHO. General guidelines for methodology on research and evaluation of traditional medicine. Geneva: World Health Organization Press; 2000a.
6. WHO. Guidelines for Appropriate Use of Herbal Medicines. Geneva: World Health Organization Press; 1998. (Accessed from on 11 August 2010) Available:<http://apps.who.int/medicinedocs/en/d/Jh2945e/11.html>
7. Cunningham AB, Shanley P, Laird S. Health, habitats and medicinal plant use. In: Colfer CJ (ed). *Human health and forests: A global overview of issues, practice and policy.* People and plants international conservation series. London: Earthscan. 2008:35–62.
8. NNMDA. The Nigeria natural medicine development agency (Federal Ministry of Science and Technology). 2008:40.
9. WHO. WHO Traditional medicine strategy 2002–2005. Geneva: WHO; 2002. (Accessed 25 October 2014) Available:[http://whqlibdoc.who.int/hq/2002/WHO EDM TRM 2002.1.pdf](http://whqlibdoc.who.int/hq/2002/WHO_EDM_TRM_2002.1.pdf),
10. World Health Organization. National policy on traditional medicine and regulation of herbal medicines. Report of a WHO global survey; 2005.
11. WHO. The legal status of traditional medicine and complimentary/alternative medicine: A worldwide review. Geneva: World Health Organization Press. 2001;16-18. (Accessed from on 11 August 2010) Available:[http://whqlibdoc.who.int/hq/2001/WHO EDM TRM 2001.2. pdf](http://whqlibdoc.who.int/hq/2001/WHO_EDM_TRM_2001.2.pdf).
12. Akunyili D. Registration and regulatory requirements for production and marketing of plant-based medicines in Nigeria: what you need to know. In: *Innovation science and bio-business development conference and Expo, herb fest at sheraton hotels and Towers Abuja*; 2003.
13. Osemeobo GJ, Ujor G. The non-wood forest products in Nigeria: report of the EC-FAO partnership programme (1998-2000). Federal department of forestry, Abuja. Forestry statistics and data collection no. AFDCA/TN/06; 1999. Available:<http://www.fao.org/docrep/003/X6695E/X6695E00.HTM>
14. Anyinam C. Availability, accessibility, acceptability, and adaptability: Four attributes of African ethno-medicine. *Soc. Sci. & Med.* 1987;25:803–11.
15. Elvin-Lewis M. Should we be concerned about herbal remedies? *J. of Ethnopharma.* 200;75(2-3):141-164.
16. Chaturvedi HK, Mahantam J, Pandey A. Treatment-seeking for febrile illness in

- north- east India: and epidemiological study for the malaria endemic zone. *Malaria J.* 2009;8:301.
17. Shaikh BT, Hatcher J. Complementary and alternative medicine in Pakistan: Prospects and limitations. Oxford: Oxford University Press; 2005.
 18. University of Maryland Medical Centre. What is Herbal Medicine?; 2010. (Accessed from on 11 October 2010. Available:<http://www.umm.edu/altmed/articles/herbal-medicine-000351.htm>)
 19. Woolf AD. Herbal remedies and children: Do they work? Are they harmful? *Pediatrics*, (2003;112(1):40-6.
 20. Abayomi Sofowora, Eyitope Ogunbodede, Adedeji Onayade. The role and place of medicinal plants in the strategies for disease prevention. *Afr. J. Tradit. Complement Altern. Med.* 2013;10(5):210-229. Available:<http://dx.doi.org/10.4314/ajtcam.v10i5.2>
 21. Sofowora A. Medicinal plants and traditional medicine in Africa. John Wiley and Sons Ltd., New York. 1984:1-20.
 22. Borris RP. Natural product research: Perspectives from a major pharmaceutical company. *J. of Ethnoph.* 1996;51:29-38.
 23. Moerman DE. An analysis of the food plants and drug plants of native North Amer. *J. of Ethnoph.* 1996;52:1-22.
 24. Ajose OA. Some Nigerian plants of dermatologic importance, *Int. J. of Derma.* 2007;46:48–55.
 25. Odugbemi T(ed.). Outlines and pictures of medicinal plants from Nigeria. 2006:1-85.
 26. African Health Monitor. Traditional medicine: Our culture, our future. A magazine of the World Health Organization Regional Office for Africa. 2003;4:1.
 27. Hoareau L. and DaSilva EJ. Medicinal plants: A re-emerging health aid. *Elect. J. of Biotech.* 1999;2:56-70.
 28. Adesina AA, Zinnah MM. Technology characteristics, farmers' perception and adoption decisions. A tobit model application in sierra leone. *Agric. Econs.* 1993;9:297-311.
 29. Goodwin B, and Mishra A. Farming efficiency and the determinants of multiple jobs holding by farm operators. *Amer. J. of Agric. Econs.* 2004;86(3):722-729.
 30. El Osta H, Ahearn M, and Mishra A. Labour supply by farm operators under decoupled farm program payments. *Review of Econs. of the Househ.* 2004;2(4):367-385.
 31. Tobit. The Use of Tobit Model in Analysing Technology Adoption: The case of Impact of Technologies on Agriculture. 1958:35-37.
 32. Gujarati DN. Basic econometrics tata McGraw hill publishing company limited. Fourth edition. New Delhi. 2004:190-278
 33. Omonona BT, Egbetokun OA, Ajjola S. and Salaam AH. Consumer preference for medicinal plants in oyo metropolis, Nigeria. *Journal of Medicinal Plants Research.* 2012;6(20):3609-3613. Available:<http://www.academicjournals.org/JMPR> DOI: 10.5897/JMPR11.1119 ISSN 1996-0875 ©2012 Academic Journals.
 34. Rikke Stamp Thorsen, Marie've Pouliot. Traditional medicine for the rich and knowledgeable: challenging assumptions about treatment-seeking behaviour in rural and periurban Nepal. *Health Poli. and Plann.* 2015;00(0):1–11. DOI: 10.1093/heapol/czv060
 35. Kroeger A. Anthropological and socio-medical health care research in developing countries. *Soci. Scien. & Med.* 1983;13B:147–61.
 36. Ahmed SM, Adams AM, Chowdhury M, Bhuiya A. Changing health seeking behaviour in Matlab, Bangladesh: Do development interventions matter? *Health Poli. and Plann.* 2000b;18:306–15.
 37. Pouliot M. Relying on nature's pharmacy in rural Burkina Faso: Empirical evidence of the determinants of traditional medicine consumption. *Soci. Scien. & Med.* 2011;73:1498-1507.
 38. Sato A. Does socio-economic status explain use of modern and traditional health care services? *Soci. Scien. & Med.* 2012b;75:1450–59.
 39. Bhattarai S, Chaudhary RP, Quave CL, Taylor RSL. The use of medicinal plants in the trans-himalayan arid zone of Mustang district, Nepal. *J. Ethnob. and Ethnom.* 2010;6:14.
 40. Shrestha PM, Dhillion SS. Medicinal plant diversity and use in the highlands of Dolakha district, Nepal. *J. of Ethnopharm.* 2003;86:81–96.
 41. Bright A. Utilization of traditional herbal medicine and its role in health care delivery in Ghana: The case of Wassa Amenfi West district. A thesis submitted to the department of geography and rural

- development, Kwame Nkrumah University of science and technology. In partial fulfilment of the requirements for the degree of master of philosophy geography and rural development faculty of social sciences, College of Art and Social Sciences. 2013; 1-122.
42. Darko IN. Ghanaian indigenous health practices: The use of herbs. Unpublished MA thesis presented to department of sociology and equity studies in education. Ontario institute for studies in education, University of Toronto; 2009.
43. Brown K. medicinal plants, indigenous medicine and biodiversity in Ghana. global environmental change working paper, 92-36, Centre for social and economic research on the global environment, University of East Anglia and University College London; 1992.
44. Falconer J, Wilson E, Asante P, Lartey J, Acquah SD, Glover EK, Beeko C, Nketiah S, Ossom K, and Lamptey M. Non-timber forest products in southern Ghana. Draft report to ODA; 1992.
45. Eisenberg DM, Kessler RC, Forster C, Norloc FE, Calkins DR, Delbanco TL. Unconventional medicine in the United States. *New England J. of Med.* 1993;328:246-252.
46. McClennon-Leong J, Ross-Kerr J. Alternative health care options in Canada. *The Canada Nurse.* 1999;11:26-30.
47. Fosu GB. Disease classification in rural Ghana: Framework and implications for health behaviour. *Soci. Scien. and Medic.* 1981;15B:471-482.
48. Rathgeber EM, Vlassoff C. Gender and tropical diseases: A new research focus. *Soci. Scien. and Med.* 1993;37(4):513-520.
49. Thorsen R. Conceptualizations of pluralistic medical fields: Exploring the therapeutic landscapes of Nepal. *Health and Place.* 2015;31:83-9.
50. Buor D. The impact of traditional medicine on health delivery services in Ghana: The Ashanti situation. *J. of the Univ. of Sci. and Tech.* 1993;13(3):138-147.

© 2018 Ariyo and Ariyo; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history/23859>*