



Comparative Physicochemical, Phytochemical Parameters and antiradical activities of *Ximenia americana* L. (Olacaceae) Almond Oils from three Locals Variants

Vincent Rouamba^a, Lassané Ouédraogo^{a,b*},
Moussa Compaoré^a, Hémayero Sama^a
and Martin Kiendrebeogo^a

^a Laboratoire de Biochimie & Chimie Appliquées, Université Joseph Ki-Zerbo, 09 BP 848 Ouagadougou 09, Burkina Faso.

^b Centre National de la Recherche Scientifique et Technologique (CNRST), Institut de l'Environnement et de Recherches Agricoles, 03 BP 7047 Ouagadougou 03, Burkina Faso.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJOCS/2023/v13i3241

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/100262>

Original Research Article

Received: 01/04/2023

Accepted: 03/06/2023

Published: 20/06/2023

ABSTRACT

Aims: This study aimed to compare local variants of *X. americana* with optimal properties for production in the food and phytomedicine industry.

Methodology: The plant material was collected from three sites in Burkina Faso, namely Gonsé, Saponé and Kokologho from May to June 2021. The oil yield was estimated and the

*Corresponding author: E-mail: lassanouedraogo.io@gmail.com;

physicochemical parameters were performed according to ISO standards. Phytochemical screening of polyphenols and flavonoids was carried out and the antioxidant activities of the oils from different trees were evaluated using DPPH and ABTS tests.

Results: The average oil extraction yield for the three variants was $42.30 \pm 14.34\%$. The estimated colour oil was yellow. The oil refraction index indicated that the Gonsé site (1.471 ± 0.002) has the highest value. The highest oil temperature value was obtained from Kokologho trees ($30.60 \pm 0.21^\circ\text{C}$) followed by the ones collected at Gonsé ($22.63 \pm 1.72^\circ\text{C}$) and Saponé ($22.60 \pm 0.41^\circ\text{C}$). The oil acid value of almonds from all three sites with an average value of 0.023 ± 0.013 mg KOH/g was lower than the international standard ISO 660-2020. The Saponé and Kokologho sites exhibited the highest oil peroxide values of 0.040 ± 0.000 mEq O₂/kg on either side, while the lowest peroxide value was found at Gonsé (0.021 ± 0.000 mEq O₂/kg). The highest saponification value is that of Kokologho with a value of 84.81 ± 4.18 mg KOH/g. The polyphenol and flavonoid content of almond oil vary within a site and between sites. Oil extracts have shown free radical scavenging capabilities (DPPH and ABTS) indicating substantial polyphenol and flavonoid content.

Conclusion: The oil has very good quality and fulfills all standard requirements for use in the phytomedicine and food industry. A mass selection can be performed within the three sites for *X. americana* plantation in agroforestry systems.

Keywords: Physicochemical; phytochemical; antioxidant activities; almond oil; *Ximania americana* L. (Olacaceae); mass selection.

1. INTRODUCTION

The occurrence of various crises known as natural disasters, conflicts and pandemics, lead decision-makers and researchers to find alternatives for the well-being of the population, especially vulnerable people [1]. Global crises sometimes affect the production chains of basic needs of the population, making them sometimes costly or even not accessible to the poorest people [2]. In recent years, the increased price of edible oils and certain oils used for therapeutic purposes could be pointed out as a result of these crises in Burkina Faso. Thus, alternative raw materials or oilseed products need to be addressed in order to overcome difficulties related to the supply of these oils. With regards to alternatives, the almonds of *Ximania americana* L. (Olacaceae) could play an important role as a source of oil in food and health because of its nutritional and therapeutic properties [3].

In fact, *Ximania americana* L. (Olacaceae) is a multi-purpose local plant in Burkina Faso used for food, pharmacopoeia, defensive hedgerows, and ornamentation. Furthermore, its fruits are used as a non-timber forest product consumed during lean seasons, but the use of the derived almonds is not well known by local communities. However; the almonds from the seeds of *X. americana* were found to contain high fat and could be used for oil and soap production [4,5]. Further studies demonstrated this oil is used in cosmetics [6,7] and treatment of skin diseases

[8]. As the oil has softening properties it can also be applied in crafts for better treatment of the leather [9].

Moreover, it was also found that oils from the almonds of *X. americana* contain high nutrients such as vitamins, protein, carbohydrates and particularly unsaturated fatty acids such as oleic and linoleic acid [10,11]. For that, the oil was recommended for nutritional use [10]. In addition, other bioactive, and phytochemicals were found [8,3]. Concerning the green fruits of *X. americana*, Almeida et al. [12] highlighted its high antioxidant activity. It can also be noted that a low content of non-essential substances was observed in the fruits, indicating that little processing would be required before it can be consumed [8].

In order to promote non-timber forest products and to find an alternative for imported oils and cosmetics, which are becoming increasingly expensive for several reasons, it becomes necessary to search for local species with high potential for vegetable oil production. For instance, *X. americana* could be one of these species. However, few studies focused on the variability of the physicochemical, phytochemical characteristics and antioxidant activities of this plant species. Therefore, this study aimed to compare local variants of *X. americana* with optimal properties for production in the food and phytomedicine industry. Our findings would contribute to improving the various uses of *X. americana* almond and also promote its

plantation as a strategy to ensure the viability of the species in its natural habitats.

2. MATERIALS AND METHODS

2.1 Study Site

The plant material was collected from three sites, namely Gonsé, Saponé and Kokologho (Fig. 1). The choice of these sites was made with regard to the evolution of *Ximenia americana* which is a wild species not yet domesticated in 3 types of contrasting environments, where the Gonsé site (highly anthropogenic classified forest in the Centre region), the site of Saponé (protected natural reserve in the South central region) and the site of Kokologho (village grove in the Central Western region of the country).

The classified forest of Gonsé is characterized by mainly ferruginous tropical sandy-clayey or gravelly soils mixed with bare soils in some places [13]. It is also a forest subject to degradation by human action. Saponé soils are gravelly, sandy, clayey and hydromorphic. The Kokologho site is a grove managed by the

villagers. On this site we find hydromorphic soils on clayey-sandy materials [14].

2.2 Collection and Sample Preparation

The identity of the plant has been confirmed by the “Herbier National du Burkina-Faso” (HNBU) located within “Centre National de Recherche Scientifique et Technologique” (CNRST) whose attribution codes are 8203N361, 8203N362 and 8203N363 for the sites of Gonsé, Saponé and Kokologho respectively.

Fruits were collected (Fig. 2) from May 28 to June 5, 2021, from eight trees (samples) selected randomly at each site according to the good condition of the tree and the presence of good fruiting. At least 500g of fruit per tree was collected. The fruit samples collected from the different sites were put in an oven at 50°C for 72 hours to eliminate the larvae. After this treatment, they were dried in the sun for a period of 2 weeks. Finally, the almonds were obtained after crushing and separating the shells. All the almond samples were then crushed using an aluminum mortar. Grinding and extraction were carried out simultaneously.




Fig. 1. Study site




Fig. 2. Shrub of *Ximenia americana*. Shrub observed in Saponé (a), Kokologho (b) and Gonsé (c). Fruit (d) and longitudinal section of the fruit (e)

2.3 Extraction

The extraction of the fat from the almond paste was carried out with hexane according to the soxhlet method for 10 hours, following the protocol of standard NF V03-924 [15].

The oil extraction yield (R) was calculated using the following equation:

$$R = \frac{W_1}{W_2} * 100$$

Where R is the extraction yield, W1 is the weight of oil extracted, and W2 is the weight of the almond paste.

2.4 Physicochemical Parameters Determination

The colour was determined using a PCE-CSM1 colorimeter. The results were expressed in terms of luminosity (L*), red-green characteristics (a*), blue-yellow characteristics (b*), chroma, relative

saturation (C*), and hue angle (H°). The refractive index, the acid value, the peroxide value and the saponification value were determined using ISO values 6320-2017, 660-2020, 3960-2017 and ISO 3657-2020 respectively.

2.5 Total Phenolic and Flavonoid Content

The total phenolic and total flavonoid contents estimation were carried out according to the method of Singleton et al. [16] adapted by Lamien-Meda et al. [17]. The polyphenol content was expressed in mg gallic acid equivalent per 1 liter (L) of extract or fraction (mg EAG/L of extract) and in mg quercetin equivalent (QE) for 1L of oil extract (mg QE/L) for total flavonoid contents.

2.6 Antiradical Evaluation Test

The evaluation of the anti-free radical activity using DPPH (2,2-diphenyl-1-picrylhydrazyl) and

ABTS ((2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) was carried out according to the method described by Chaves et al. [18], and Re et al. [19] with some modifications. The antiradical activity of the oil concentration which scavenges 50% of the free radical DPPH•+, and reduces the radical cation ABTS+• was expressed in mg/mL oil, and mg Trolox equivalent per ml (μmol TE/ml) of extract successively.

2.7 Statistical Analysis

The data were analyzed with the Software Minitab version 19.1. An analysis of variance (ANOVA) was carried out in order to detect any significant differences between the trees or the sites for the characters studied. The separation of the means was performed by Tukey's test at the 5% threshold.

3. RESULTS AND DISCUSSION

3.1 Extraction Yield

The extraction yields obtained vary from $36.88 \pm 9.17\%$; $42.12 \pm 14.85\%$ and $47.92 \pm 10.02\%$ for the sites of Gonsé, Saponé and Kokologho, respectively. The average yield for all three sites is 42.30 ± 14.34 . There is no significant difference ($P > 0.05$) between the yields obtained in the three sites. These yields are lower than the yields obtained by Oladipo, et al. [20] and Satoto et al. [6] using *X. americana* samples from Nigeria and Angola, which were 45.7% and 61.78%, respectively. There are multiple factors that could explain these differences: different pedoclimatic conditions, differences in extraction solvents and the conditions for collection, drying and conservation of the almonds used by these authors.

3.2 Physicochemical Parameters

The physicochemical parameters per site are summarized in Table 1 (colour) and Table 2 (refractive index, acid value, peroxide value and saponification value), while the comparison of physicochemical parameters between the sites is present in Fig. 3.

The colour angles of the oil extracted from the various trees of Gonsé site vary between 78.90 and 82.33 corresponding to the colour index PY 108 (anthrapyrimidine yellow) to PY35 (cadmium

yellow). The hue angles of Saponé trees samples vary from 77.87 to 86.46 corresponding to the colour index PY108 to the colour index PY129 (green gold). The hue angles for Kokologho trees samples vary from 70.08 to 89.90 corresponding to the colour index PY1 (hansa yellow G) to the colour index PY138 (quinophthalone yellow). In general, we can say that the colour of the oil is yellow. For example, the work of Hozaifa Medaowe Ahmed [21] showed that the oil of *X. americana* has a bright yellow colour that is line with our results.

Comparing the refractive index between the three study sites, a significant difference ($p \leq 0.001$) between the Gonsé, Saponé and Kokologho sites can be observed. The Gonsé site, particularly trees 1 and 2 indicate the highest value of about 1.47. The refractive index values obtained from all our samples were higher than those found by Tanko et al. [22] in Nigeria which was 1.413. These variations could be explained by the temperatures of the oils but also by the extent of refining of the oils. Some authors such as Oladipo et al. [20], and Sharief et al. [10] who also worked on *X. americana* oils found refractive index values similar to ours (1.467 and 1.470). According to the ISO 6320-2017 standard, the values obtained in our study indicate a good purity of the oils. The refractive indices of the almonds harvested from all our study sites demonstrate that the almond oil of *X. americana* could be used as a food oil.

The highest oil temperature value was found in the site of Kokologho for all the tree samples (about 30°C) with a significant difference ($P < 0.001$) regarding the other site's trees samples.

In Gonsé, the minimum and maximum temperature of the almond oils was respectively $21.13 \pm 0.15^\circ\text{C}$ (tree 4) and $26.70 \pm 0.85^\circ\text{C}$ (tree 1). At Saponé the minimum was $21.93 \pm 0.058^\circ\text{C}$ (tree 8) and the maximum was $23.23 \pm 0.15^\circ\text{C}$ (tree 5).

The highest acid value in Gonsé is 0.07 ± 0.00 mg KOH/g (tree 6). At Saponé the highest acid value (0.1 ± 0.00 mg KOH/g) is from tree 2 and the smallest value (0.007 ± 0.00 mg KOH/g) derives from tree 3, while at the Kokologho site, the highest value is 0.05 ± 0.00 (tree 8) and the smallest (0.002 ± 0.00 mg KOH/g) is that from tree 5.

Table 1. Colour coordinates (CIELab) of *Ximenia americana* almond oils per site

Sites	Trees	L*	a*	b*	C*	H°
Gonsé	Tree ₁	47.21 ± 3.01 ^a	1.96 ± 0.14 ^a	14.93 ± 3.21 ^a	15.20 ± 3.21 ^a	78.95 ± 2.66 ^a
	Tree ₂	47.20 ± 3.00 ^a	1.92 ± 0.19 ^a	14.92 ± 3.21 ^a	15.19 ± 3.21 ^a	78.92 ± 2.67 ^a
	Tree ₃	47.19 ± 3.00 ^a	1.91 ± 0.19 ^a	14.89 ± 3.19 ^a	15.18 ± 3.21 ^a	78.91 ± 2.67 ^a
	Tree ₄	47.18 ± 3.01 ^a	1.76 ± 0.22 ^a	14.88 ± 0.78 ^a	15.17 ± 3.21 ^a	78.90 ± 2.67 ^a
	Tree ₅	47.18 ± 3.01 ^a	2.90 ± 0.78 ^a	12.50 ± 0.78 ^a	12.63 ± 0.80 ^a	82.33 ± 1.31 ^a
	Tree ₆	47.19 ± 3.01 ^a	2.91 ± 0.78 ^a	12.51 ± 0.80 ^a	12.64 ± 0.80 ^a	82.34 ± 1.31 ^a
	Tree ₇	49.03 ± 2.06 ^a	1.91 ± 0.40 ^a	13.30 ± 1.74 ^a	13.43 ± 1.75 ^a	81.84 ± 1.57 ^a
	Tree ₈	49.02 ± 2.06 ^a	1.90 ± 0.40 ^a	13.30 ± 1.74 ^a	15.20 ± 1.75 ^a	81.84 ± 1.57 ^a
Saponé	Tree ₁	62.96 ± 5.78 ^a	1.23 ± 0.51 ^b	22.51 ± 1.74 ^a	22.55 ± 1.75 ^a	86.44 ± 1.60 ^a
	Tree ₂	63.03 ± 5.75 ^a	1.28 ± 0.52 ^b	22.54 ± 1.76 ^a	22.58 ± 1.73 ^a	86.46 ± 1.60 ^a
	Tree ₃	47.29 ± 0.90 ^b	2.36 ± 0.28 ^{ab}	11.76 ± 0.30 ^b	12.01 ± 0.31 ^b	79.79 ± 1.21 ^b
	Tree ₄	47.31 ± 0.91 ^b	2.37 ± 0.28 ^{ab}	11.80 ± 0.33 ^b	12.02 ± 0.31 ^b	79.80 ± 1.21 ^b
	Tree ₅	47.40 ± 1.00 ^b	2.40 ± 0.30 ^{ab}	11.81 ± 0.33 ^b	12.03 ± 0.30 ^b	79.83 ± 1.20 ^b
	Tree ₆	52.20 ± 1.08 ^b	3.13 ± 0.62 ^{ab}	14.51 ± 2.16 ^b	14.85 ± 2.22 ^b	77.87 ± 1.26 ^b
	Tree ₇	52.21 ± 1.08 ^b	3.14 ± 0.62 ^a	14.52 ± 2.16 ^b	14.88 ± 2.21 ^b	77.87 ± 1.26 ^b
	Tree ₈	52.22 ± 1.08 ^b	3.15 ± 0.62 ^a	14.53 ± 2.60 ^b	14.88 ± 2.21 ^b	77.96 ± 1.22 ^b
Kokologho	Tree ₁	52.40 ± 4.90 ^b	2.02 ± 0.86 ^{ab}	19.21 ± 1.38 ^a	19.44 ± 1.22 ^a	83.90 ± 2.84 ^a
	Tree ₂	41.82 ± 1.65 ^b	3.54 ± 0.28 ^a	09.76 ± 0.43 ^b	10.38 ± 0.50 ^b	70.08 ± 0.71 ^b
	Tree ₃	52.40 ± 4.90 ^b	2.02 ± 0.86 ^{ab}	19.21 ± 1.38 ^a	19.44 ± 1.22 ^a	83.90 ± 2.84 ^a
	Tree ₄	52.40 ± 4.90 ^b	2.06 ± 0.86 ^{ab}	19.21 ± 1.38 ^a	19.44 ± 0.90 ^a	83.90 ± 2.84 ^a
	Tree ₅	41.82 ± 1.65 ^b	3.54 ± 0.28 ^a	09.76 ± 0.43 ^b	10.38 ± 0.50 ^b	70.08 ± 0.71 ^b
	Tree ₆	52.40 ± 4.90 ^b	2.02 ± 0.86 ^{ab}	19.21 ± 1.37 ^a	19.44 ± 1.22 ^a	83.90 ± 2.84 ^a
	Tree ₇	74.73 ± 7.77 ^a	0.00 ± 0.00 ^b	19.27 ± 0.89 ^a	19.30 ± 0.90 ^a	89.90 ± 3.70 ^a
	Tree ₈	74.73 ± 7.77 ^a	0.03 ± 0.00 ^b	19.27 ± 0.89 ^a	19.30 ± 0.90 ^a	89.90 ± 3.70 ^a

Results are presented as mean ± SD (n=3). Values that do not share any letter are significantly different.
 Legend: L*: Luminosity, a*: Red-green characteristics, b*: Blue-yellow characteristics, C*: Chroma. H°: Hue angle.

Table 2. Physicochemical parameters of *Ximenia americana* almond oil per site

Sites	Trees	RI	AV	PV	SV	T (°C)
Gonsé	Tree ₁	1.470 ± 0.00 ^a	0.06 ± 0.02 ^{ab}	0.03 ± 0.01 ^a	79.31 ± 5.68 ^a	26.70 ± 0.85 ^a
	Tree ₂	1.473 ± 0.00 ^a	0.04 ± 0.00 ^b	0.02 ± 0.00 ^b	79.31 ± 5.68 ^a	22.43 ± 0.50 ^b
	Tree ₃	1.471 ± 0.00 ^a	0.04 ± 0.00 ^b	0.02 ± 0.00 ^b	53.29 ± 8.42 ^b	21.43 ± 0.15 ^{bc}
	Tree ₄	1.470 ± 0.00 ^a	0.04 ± 0.00 ^b	0.02 ± 0.00 ^b	81.02 ± 2.79 ^a	21.13 ± 0.15 ^c
	Tree ₅	1.470 ± 0.00 ^a	0.04 ± 0.00 ^b	0.02 ± 0.00 ^b	86.77 ± 8.23 ^a	22.50 ± 0.10 ^b
	Tree ₆	1.471 ± 0.00 ^a	0.07 ± 0.00 ^a	0.02 ± 0.00 ^b	89.58 ± 0.18 ^a	22.30 ± 0.10 ^b
	Tree ₇	1.474 ± 0.00 ^a	0.04 ± 0.00 ^b	0.02 ± 0.00 ^b	94.61 ± 0.00 ^a	22.10 ± 0.36 ^{bc}
	Tree ₈	1.469 ± 0.00 ^a	0.04 ± 0.00 ^b	0.02 ± 0.00 ^b	89.59 ± 5.78 ^a	22.43 ± 0.06 ^b
Saponé	Tree ₁	1.470 ± 0.00 ^{ab}	0.004 ± 0.00 ^f	0.04 ± 0.00 ^a	70.12 ± 2.8 ^c	22.43 ± 0.06 ^{cd}
	Tree ₂	1.473 ± 0.00 ^a	0.049 ± 0.00 ^a	0.04 ± 0.00 ^a	81.34 ± 2.8 ^a	22.70 ± 0.10 ^{bc}
	Tree ₃	1.470 ± 0.00 ^{ab}	0.007 ± 0.00 ^d	0.04 ± 0.00 ^a	78.54 ± 0.00 ^{ab}	22.63 ± 0.06 ^{bc}
	Tree ₄	1.468 ± 0.00 ^{ab}	0.008 ± 0.00 ^c	0.04 ± 0.00 ^a	84.15 ± 0.00 ^a	22.93 ± 0.15 ^{ab}
	Tree ₅	1.470 ± 0.00 ^{ab}	0.008 ± 0.00 ^c	0.04 ± 0.00 ^a	81.34 ± 2.8 ^a	23.23 ± 0.15 ^a
	Tree ₆	1.470 ± 0.00 ^{ab}	0.030 ± 0.00 ^b	0.04 ± 0.00 ^a	81.34 ± 2.81 ^a	22.73 ± 0.12 ^{bc}
	Tree ₇	1.463 ± 0.00 ^b	0.030 ± 0.00 ^b	0.04 ± 0.00 ^a	84.15 ± 0.00 ^a	22.20 ± 0.30 ^{de}
	Tree ₈	1.466 ± 0.00 ^{ab}	0.006 ± 0.00 ^e	0.04 ± 0.00 ^a	72.93 ± 0.00 ^{bc}	21.93 ± 0.06 ^e
Kokologho	Tree ₁	1.463 ± 0.00 ^a	0.008 ± 0.00 ^d	0.04 ± 0.00 ^b	92.56 ± 2.80 ^a	30.60 ± 0.00 ^c
	Tree ₂	1.465 ± 0.00 ^a	0.004 ± 0.00 ^{ef}	0.04 ± 0.00 ^b	83.81 ± 5.59 ^{ab}	30.73 ± 0.06 ^b
	Tree ₃	1.467 ± 0.00 ^a	0.020 ± 0.00 ^c	0.04 ± 0.00 ^b	86.95 ± 2.81 ^{ab}	30.93 ± 0.06 ^a
	Tree ₄	1.462 ± 0.00 ^a	0.030 ± 0.00 ^b	0.04 ± 0.00 ^b	78.54 ± 5.61 ^b	30.80 ± 0.00 ^b
	Tree ₅	1.465 ± 0.00 ^a	0.002 ± 0.00 ^f	0.04 ± 0.00 ^b	84.15 ± 5.61 ^{ab}	30.60 ± 0.00 ^c
	Tree ₆	1.465 ± 0.00 ^a	0.015 ± 0.00 ^c	0.04 ± 0.00 ^b	81.34 ± 8.42 ^{ab}	30.40 ± 0.00 ^{de}
	Tree ₇	1.465 ± 0.00 ^a	0.005 ± 0.00 ^e	0.04 ± 0.00 ^b	84.15 ± 0.00 ^{ab}	30.46 ± 0.06 ^d
	Tree ₈	1.467 ± 0.00 ^a	0.050 ± 0.00 ^a	0.06 ± 0.02 ^a	86.95 ± 2.80 ^{ab}	30.30 ± 0.00 ^e

Results are presented as mean ± SD (n=3). Values that do not share any letter are significantly different.

Legend: RI: Refraction index; AV: Acid value (mg KOH/g); PV: Peroxide value (mEq O₂/kg); SV: Saponification value (mg KOH/g), T (°C): Temperature in degrees Celsius, 1 to 8: Tree number




Fig. 3. Comparison of *Ximenia americana* almond oil physicochemical parameters between the sites: acid value (a), peroxide value (b), refractive index (c), saponification value (d), temperature (e)

The comparison between the acid values of the three sites shows that the Gonsé sample from tree 6 has the highest value and the lowest is from tree 5 of Kokologho. The acid values are lower than the international standard ISO 660-2020 for fatty substances where for refined oils the limit is 0.6 mg KOH/g oil, while that of virgin fats and oils can reach an acid value of 4.0 mg KOH/g fat or oil [23]. Therefore, according to the Codex Alimentarius *X. americana* oil can be considered as food. Authors such Eromosele et al. [24] and [22] found values of 0.14 mg KOH/g, 0.29 ± 0.15 mg KOH/g, respectively, which are much higher than our values. On the other hand, some authors such as Oladipo et al. [20] obtained a value of 16.13 mg KOH/g and which was much higher than the international standard. This very low content of free fatty acids in our oil extracts derived from *X. americana* seeds is an indication that it will have good long-term stability.

The peroxide value of tree 1 (0.03 ± 0.01 mEq O₂/kg) in the Gonsé site is significantly different

($P \leq 0.001$) from the rest of the trees sampled. For the Kokologho and Saponé sites, the peroxide values are the same (0.040 ± 0.00 mEq O₂/kg) for all trees.

In general, the values found in our samples were lower than those obtained by Saeed and Bashier [25] and Sharief et al. [10] which were 30 mEq O₂/kg oil from *X. americana*. These low values of peroxide could derive either from the presence of antioxidants (polyphenols, flavonoids, etc.) in our oil samples, which prevent the oil from going rancid. Also, suboptimal storage conditions (temperature, light) could increase oxidation processes in crude oil. Moreover, non-similarity in the sample processing could be the main reason for these differences. As the Codex Alimentarius standards [23] set the peroxide value for refined oils at a value less than or equal to 10 mEqO₂/kg and for crude oils at 15 mEqO₂/kg, our values are far below these limits.

The saponification values were significantly different ($P \leq 0.001$) between trees of the same

site for all locations studied. In Gonsé, tree 7 (94.61 ± 0.00 mg KOH/g) has the highest saponification value and the lowest comes from tree 3 (53.29 ± 8.40 mg KOH/g). At Saponé, trees 4 and 7 show the highest values (84.15 ± 0.00 mg KOH/g) and the lowest saponification value is from tree 1 (70.12 ± 2.8 mg KOH/g). As for the Kokologho site, the saponification value of tree 1 (92.56 ± 2.80 mg KOH/g) was the highest and the lowest value was that of tree 4 (78.54 ± 5.61 mg KOH/g).

The saponification values obtained were lower than those described by the authors Satoto et al. [6] and Tanko et al. [22] which were respectively 295.00 ± 0.17 mg KOH/g and 179.94 ± 1.69 mg KOH/g oil. However, other authors like Saeed & Bashier [25] found values of 11.43 mg KOH/g which were much lower than our results. These differences could be due either to non-similarity sample processing or the fact that our oils contain less free fatty acids, or to different climatic, pedoclimatic or edaphic conditions. In addition, poor separation of the soap during neutralization or by poor washing [26] could be a reason for the differences found. Our low saponification values also indicate that our oils contain a long chain of fatty acids as could be found in other vegetable oils (such as sunflower) or animal oils (such as fatty chicken) [27].

3.3 Total Polyphenol and Flavonoid Content in *Ximenia americana* Almonds Oil

The total polyphenol and flavonoid content per site are indicated in Table 3 and the comparison between the sites is mentioned in Fig. 4a and Fig. 4b. Within each site, there is a significant difference ($p \leq 0.001$) between trees for the total polyphenol content in *X. americana* almond oils. In Gonsé, tree 1 (346.02 ± 13.43 mg EAG/L) and tree 2 (339.8 ± 47.5 mg EAG/L) present the highest values of total polyphenols and the lowest value is from tree 3 (143.02 ± 12.52 mg EAG/L). At Saponé the polyphenol content of almond oil from tree 3 (441.98 ± 0.606 mg EAG/L) is significantly different from the other 7 trees studied.

At Kokologho, the content is 321.37 ± 10.1 mg EAG/L for tree 8 and is statistically different from either other trees. The values of the total polyphenol content in the oils of *X. americana* compared between the sites indicate that the results are statistically different ($p \leq 0.001$) from each other, but the highest content is from saponé tree 3 (442 ± 0.60 mg EAG/L).

The total polyphenol content variation is high within the same site, this can be due to the state of the tree (biotic and abiotic factors) in its native environment. In the same way, Laitinen et al. [28] were underlined in their work entitled variation in phenolic compounds within a birch population that high chemical variation may be an effective means of adaptation in variable abiotic and biotic environments of trees.

In Gonsé site, tree 1 (34.76 ± 1.69 mg QE/L) presented a high flavonoid content compared to the other trees. At Saponé, the flavonoid content of almond oil (16.17 ± 0.17 mg QE/L) is significantly different ($p \leq 0.001$) for tree 6 compared to the remaining trees. In Kokologho, tree 7 with a content of 37.57 ± 4.22 mg QE/L is statistically different from the other trees.

When the 3 sites are taken into account together, the total flavonoid content of the Kokologho tree 7 is the highest among the remaining trees.

3.4 Antiradical Activities

The concentrations of *X. americana* tree almond oil extracts from each site that inhibited 50% of the DPPH radical (IC₅₀), and ABTS radical reduction activities are presented in Table 3. Analyzes of IC₅₀ values of *X. americana* tree almond oils within each site showed highly significant variability ($p \leq 0.001$) when individual trees are compared.

For the Gonsé site, the lowest 50% inhibition value could be detected in the oil fractions from tree 4 (0.11 ± 0.03 mg/mL) and tree 3 (0.14 ± 0.01 mg/mL). Concerning the Saponé site, the minimum oil concentration to inhibit 50% of the DPPH radical is from tree 1 (0.11 ± 0.04 mg/mL). As for the Kokologho site, the lowest IC₅₀ value is for tree 5 (0.09 ± 0.02 mg/mL). The oil fraction from this tree has the highest antioxidant activity of all trees tested.

There is a significant difference between the IC₅₀ values of the three sites ($P \leq 0.001$) (Fig. 4c).

Taken together, all *X. americana* oil extracts tested for DPPH scavenging activities show radical scavenging capabilities. While Bazezew et al. [7] reported that the oil from the seed of *X. americana* had radical scavenging capabilities indicating that the seeds of *X. americana* had a capacity to scavenge DPPH* radicals at a concentration of 200 µg/mL and an IC₅₀ of 154 µg/ml.

Table 3. Total polyphenolic, contents, antiradical activities per site

Sites	Trees	TPC	TFC	Anti-DPPH	Anti-ABTS
Gonsé	Tree ₁	346.02 ± 13.43 ^a	34.76 ± 1.69 ^a	0.63 ± 0.06 ^a	32.75 ± 0.43 ^b
	Tree ₂	339.80 ± 47.50 ^a	28.32 ± 1.56 ^b	0.50 ± 0.29 ^{ab}	31.80 ± 0.35 ^c
	Tree ₃	143.02 ± 12.52 ^d	3.90 ± 0.13 ^d	0.14 ± 0.01 ^c	33.58 ± 0.06 ^a
	Tree ₄	241.07 ± 6.50 ^b	1.15 ± 0.05 ^d	0.11 ± 0.03 ^c	32.10 ± 0.63 ^c
	Tree ₅	178.60 ± 25.90 ^{cd}	29.56 ± 2.13 ^b	0.20 ± 0.14 ^{bc}	31.41 ± 0.04 ^c
	Tree ₆	254.61 ± 4.65 ^b	18.53 ± 1.05 ^b	0.40 ± 0.02 ^{abc}	30.30 ± 0.76 ^d
	Tree ₇	238.19 ± 1.13 ^{bc}	30.97 ± 1.36 ^b	0.40 ± 0.01 ^{abc}	28.73 ± 0.28 ^e
	Tree ₈	202.60 ± 17.50 ^{bcd}	30.94 ± 0.36 ^b	0.40 ± 0.05 ^{abc}	33.45 ± 0.44 ^a
Saponé	Tree ₁	51.07 ± 0.00 ^f	9.80 ± 0.13 ^b	0.14 ± 0.04 ^c	34.31 ± 0.34 ^{abc}
	Tree ₂	197.67 ± 2.52 ^{cd}	6.81 ± 0.17 ^d	0.16 ± 0.06 ^c	34.21 ± 0.10 ^a
	Tree ₃	441.98 ± 0.60 ^a	8.56 ± 0.05 ^c	0.27 ± 0.10 ^{bc}	30.52 ± 0.63 ^{bc}
	Tree ₄	144.40 ± 0.40 ^e	6.15 ± 0.05 ^e	0.24 ± 0.00 ^{bc}	30.52 ± 1.04 ^c
	Tree ₅	206.17 ± 1.53 ^c	4.312 ± 0.09 ^f	0.45 ± 0.03 ^a	26.67 ± 2.03 ^d
	Tree ₆	282.05 ± 9.64 ^b	16.17 ± 0.17 ^a	0.16 ± 0.00 ^c	23.19 ± 0.66 ^e
	Tree ₇	286.40 ± 23.30 ^b	4.12 ± 0.09 ^f	0.33 ± 0.00 ^{ab}	26.68 ± 1.39 ^d
	Tree ₈	178.95 ± 4.65 ^d	6.04 ± 0.09 ^e	0.24 ± 0.00 ^{bc}	23.63 ± 0.03 ^d
Kokologho	Tree ₁	196.22 ± 14.95 ^{cd}	29.02 ± 1.55 ^b	0.51 ± 0.00 ^a	26.55 ± 0.83 ^d
	Tree ₂	67.64 ± 4.04 ^g	9.85 ± 0.43 ^c	0.51 ± 0.06 ^a	29.81 ± 0.50 ^{ab}
	Tree ₃	254.10 ± 12.32 ^b	26.34 ± 0.82 ^b	0.53 ± 0.07 ^a	28.08 ± 0.33 ^{cd}
	Tree ₄	171.88 ± 12.53 ^{de}	4.00 ± 0.55 ^d	0.36 ± 0.04 ^b	23.79 ± 0.62 ^e
	Tree ₅	130.80 ± 10.26 ^f	7.24 ± 1.12 ^{cd}	0.08 ± 0.02 ^d	29.16 ± 0.50 ^{bc}
	Tree ₆	208.44 ± 4.24 ^c	7.85 ± 1.72 ^{cd}	0.20 ± 0.00 ^c	30.76 ± 0.68 ^{ab}
	Tree ₇	149.66 ± 1.61 ^{ef}	37.57 ± 4.22 ^a	0.20 ± 0.00 ^{cd}	27.53 ± 0.19 ^{cd}
	Tree ₈	321.37 ± 10.10 ^a	8.76 ± 0.55 ^{cd}	0.11 ± 0.02 ^{cd}	30.84 ± 0.39 ^a

Results are presented as mean ± SD (n=3). Values that do not share any letter are significantly different.

Legend: TPC: Total polyphenol content (mg EAG/L, oil), TFC: Total flavonoid content (mg QE/L, oil), Anti-DPPH (mg/mL), Anti-ABTS (μmol TE /mL, oil), 1 to 8: Tree number.




Fig. 4. Comparison of *Ximenia americana* almond oil total polyphenolic, contents, antiradical activities between the sites: Total polyphenol content (a), Total flavonoid content (b), anti-DPPH (c), Anti-ABTS (d)

Work carried out on the DPPH antiradical activity of the fresh fruits of *X. americana* had shown DPPH* radical scavenging capacities at 68.56 ± 0.10 mg TE/100 g [29]. But all this author used a completely different processing procedure.

At Gonsé the ABTS radical reduction activities of the almond oils of tree 3 (33.60 ± 0.06) μ mol Trolox equivalent /ml) and 8 (33.45 ± 0.44 μ mol Trolox equivalent /ml) are significantly different from trees 1, 2, 4, 5, 6 and 7. At Saponé, tree 1 (34.59 ± 0.34 μ mol Trolox equivalent /ml) and 2 (34.21 ± 0.10 μ mol Trolox equivalent /ml) present significantly high activities compared to the other trees. At Kokologho, it is the oils from trees 6 (30.76 ± 0.68 μ mol Trolox equivalent /ml) and 8 (30.84 ± 0.39) which presented a more significant activity compared to the other trees.

The between-site comparison of the average antioxidant activities of almond oils used for the ABTS test showed a significant difference (Fig. 4d).

The variation of the antioxidant activity within the same site could be explained by the disparity of the trees in terms of morphology or physiology or by an ongoing defense reaction against bio-aggressors.

These results obtained by site demonstrate that our samples contain molecules capable of donating electrons (or hydrogen).

Taken together the nutritional potential of *X. americana* almond oil could be substantiated by the presence of polyphenols and flavonoid antioxidant activities. Some studies have pointed out that the seeds of *X.americana* contain hydrocyanic acid but the toxicity of the oil has not been proven [30]. Moreover, the oil from the seed is used in food, cosmetics (for hair and skin) and is marketed in Ethiopia [3]. These same authors and other authors like Kefelegn and Desta [3] affirm that if the plant is studied it would have many advantages for food safety, for medicine and could be a basic raw material in the food industry.

4. CONCLUSION

The physicochemical parameters, noted as acid, peroxide and saponification values of *X. americana* oils from the three locations were within the threshold values proposed by the Codex Alimentarius. The parameters of *X. americana* almond oils varied somewhat from site to site. The extraction yield of the oils was favorable and was slightly high with the almonds from the Kokologho site compared to the other sites.

In addition, phytochemical screening of the almond oil from *X. americana* showed significant content of phenolic compounds and flavonoids

providing the oil with substantial free radical scavenging capacities (DPPH and ABTS).

This study indicates, that the oil of *X. americana* could be used in phytomedicine and food processing. In order to promote and preserve the natural populations of this multipurpose species in Burkina Faso, we propose to make a mass selection of this species for plantation.

DECLARATION

This study does not involve experiments on animals or human subjects.

ACKNOWLEDGEMENTS

The authors sincerely thank Josef Endl, Weilheim, Germany, and Salam Sawadogo, CNRST, Ouagadougou, Burkina Faso for many helpful corrections and recommendations.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Machlis GE. Science during crisis: The role of science in disaster response. WIT Transactions on the Built Environment. 2019;190:103-111.
2. FAO. State of food insecurity in the world 2009: Economic crises-impacts and lessons learned. Rome:World Health Organization: Food and Agriculture Organization of the United Nations; 2009.
3. Kefelegn GA, Desta B. *Ximenia americana*: Economic Importance, Medicinal Value, and Current Status in Ethiopia. The Scientific World Journal. 2021;2021:8880021. Available:https://doi.org/10.1155/2021/8880021
4. Vermaak I, Kamatou G, Komane-mofokeng B, Viljoen AM, Beckett K. African seed oils of commercial importance - Cosmetic applications. South African Journal of Botany. 2011; 77(4):920-933. Available:http://dx.doi.org/10.1016/j.sajb.2011.07.003
5. Urso V, Signorini MA, Bruschi P. Survey of the ethnobotanical uses of *Ximenia americana* L. (mumpeke) among rural communities in South Angola. Journal of Medicinal Plants Research. 2013;7(1):7-18.
6. Satoto G, Fernandes AS, Saraiva N, Santos F, Neng N, Nogueira M, et al. An Overview on the Properties of *Ximenia* Oil Used as Cosmetic in Angola. Biomolecules. 2019;10(1):18-29. Available:http://dx.doi.org/10.3390/biom10010018
7. Bazezew AM, Emire SA, Sisay MT. Bioactive composition, free radical scavenging and fatty acid profile of *Ximenia americana* grown in Ethiopia. Heliyon. 2021;7:e07187. Available:https://doi.org/10.1016/j.heliyon.2021.e07187. Available:http://dx.doi.org/10.5897/JMPR11.492
8. Muhammad A, Haruna SY, Birnin-yauri AU, Muhammad AH, Elinge CM. Nutritional and anti-nutritional composition of *Ximenia americana* fruit. American Journal of Applied Chemistry. 2019;7(4): 123–129. Available:http://dx.doi.org/10.11648/j.ajac.20190704.13
9. Arbonnier M. Arbres, arbustes et lianes des zones sèches d'Afrique de l'Ouest. Montpellier: CIRAD; MNHN; 2000.
10. Sharief TM, Mohammed Bashier RS, Haroon MI. Phytochemical evaluation and uses of *Ximenia americana* L. in Central Darfur. International Journal of Current Microbiology and Applied Sciences. 2022;11(2):353-360. Available:https://doi.org/10.20546/ijcmas.2022.1102.040
11. Kundlik PS, Rajasekaran S. Ethnopharmacological review of *Ximenia americana* and *Lindera communis*. European Journal of Molecular and Clinical Medicine. 2023;10(1):4370-4381.
12. Almeida MLB, Freitas WEDS, Morais PLD De, Sarmiento JDA, Alves RE. Bioactive compounds and antioxidant potential fruit of *Ximenia americana* L. Food Chemistry. 2016;192:1078–1082. Available:http://dx.doi.org/10.1016/j.foodchem.2015.07.129
13. Sankara BT. Climate variability and natural resource management: Case of the classified forest and partial wildlife reserve of Gonse in Burkina Faso. Master Thesis, Université de Ouagadougou, Burkina Faso; 2010.

14. Sawadogo O. Notice of environmental and social impact of the project to build simplified drinking water supply systems in the localities of Gourcy (municipality of Nandiala), Ouoro (Sourgou), Sakoinzé (Kokologho); province of Boulkiemdé, Center Region.
Accessed 20 December 2022.
Available: https://www.afdb.org/sites/default/files/nies_aeps_pasepa-2r_sanguie_vf.pdf
15. Novidzro KM, Wokpor K, Fagla BA, Koudouvo K, Dotse K, Osseyi E, et al. Study of some physicochemical parameters and analysis of mineral elements, chlorophyll pigments and carotenoids of *Griffonia simplicifolia* seed oil. *International Journal of Biological and Chemical Sciences*. 2019;13(4):2360-2373.
Available: <https://doi.org/10.4314/ijbcs.v13i4.38>
16. Singleton VL, Orthofer R, Lamuela-Raventos RM. Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Methods Enzymol*. 1999;299:152-78.
17. Lamien-Meda A, Lamien EC, Compaoré MY, Meda, Roland NT, Kiendrebeogo M, Zeba B, et al. Polyphenol content and antioxidant activity of fourteen wild edible fruits from Burkina Faso. *Molecules*. 2008;13(3):581-594.
Available: <https://doi.org/10.3390/molecules13030581>
18. Chaves N, Santiago A, Alías JC. Quantification of the antioxidant activity of plant extracts: Analysis of sensitivity and hierarchization based on the method used. *Antioxidant*. 2020;9:76-90.
19. Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radical Biology and Medicine*. 1999;26(9/10):1231-1237.
20. Oladipo GO, Eromosele IC, Folarin OM. Formation and characterization of paint based on alkyd resin derivative of *Ximenia americana* (Wild Olive) seed oil. *Environment and Natural Resources Research*. 2013;3(3):52-62.
Available: <https://doi.org/10.5539/enrr.v3n3p52>
21. Hozaiifa Medaowe Ahmed A. Physico-chemical properties, gas chromatography-mass spectrometric analysis and antioxidant activity of Um Wedeka (*Ximenia americana* . L) seed oil. Master thesis, University of Gezira, Sudan; 2020.
22. Tanko E, Ajai AI, Lafiya RA, Dauda BEN, Mathew JT, Omozokpia J. Physico-chemical , fatty acid profile and amino acid composition of the fruit pulp and seeds of *Ximenia americana* L . (Tallow Plum) obtained in Niger State , Nigeria. *International Journal of Food Chemistry*. 2017;1(2):30-34.
Available: <http://dx.doi.org/10.11648/j.ijfc.20170102.11>
23. Joint FAO/WHO Codex Alimentarius Commission. *Codex Alimentarius: Food hygiene, basic texts*. Rome :World Health Organization : Food and Agriculture Organization of the United Nations; 2003.
24. Eromosele IC, Eromosele CO, Akintoye AO, Komolafe TO. Characterization of oils and chemical analyses of the seeds of wild plants. *Plant Foods for Human Nutrition*. 1994;46:361–365.
Available: <https://doi.org/10.1007/BF01088437>
25. Saeed AEM, Bashier RSM. Physico-chemical analysis of *Ximenia americana* . L seed oil and structure elucidation of some chemical constituents of its seed oil and fruit pulp. *Journal of Pharmacognosy and Phytotherapy*. 2010;2(4):49-55.
Available: <https://doi.org/10.5897/JPP.9000042>
26. Diakite K, Diagouraga S, Diawara M. Study of the physicochemical parameters of cotton seed oils produced in the CMDT zone in Mali. *International Journal of Biological and Chemical Sciences*. 2022;16(3):1320–1330.
Available: <https://doi.org/10.4314/ijbcs.v16i3.33>
27. Ivanova M, Hanganu A, Dumitriu R, Tociu M, Ivanov G, Stavarache C, et al. Saponification value of fats and oils as determined from data: The case of dairy fats. *Foods*. 2022;11:1466-1473.
Available: <https://doi.org/10.3390/foods1101466>
28. Laitinen ML, Julkunen-Tiitto R, Rousi M. Variation in phenolic compounds within a birch (*Betula pendula*) population. *Journal of Chemical Ecology*. 2000;26:1609–1622.
Available: <https://doi.org/10.1023/A:1005582611863> .

29. Koumba BB, Bayang JP, Wangso H, Kolla MC, Laya A. Variation of phenolics (Bound and Free), minerals, and antioxidant activity of twenty-eight wild edible fruits of twenty-three species from Far North Region of Cameroon. *BioMed Research International*. 2021; 2021:14. Available: <https://doi.org/10.1155/2021/4154381>.
30. Maiga A, Diallo D, Fane S, Sanogo R, Smestad B, Cisse B. A survey of toxic plants on the market in the district of Bamako, Mali: Traditional knowledge compared with a literature search of modern pharmacology and toxicology. *Journal of Ethnopharmacology*. 2005;96: 183–193. Available: <https://doi.org/10.1016/j.jep.2004.09.005>

© 2023 Rouamba et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/100262>