



## Difference of Propolis Administration as an Antioxidant against Oxidative Stress (Malondialdehyde Level (MDA)) and Apoptosis (Caspase-3 Expression) in Skin Graft Model White Rats (*Rattus norvegicus*)

Ricky Dwi Nur Tyastono<sup>1\*</sup>, Amru Sungkar<sup>1</sup>, Kristanto Yuli Yarso<sup>1</sup>, Dewi Haryanti<sup>1</sup>, Nunik Agustrina<sup>1</sup>, Ida Bagus Budi<sup>1</sup> and Brian Wasita<sup>2</sup>

<sup>1</sup>Department of Surgery, Dr. Moewardi Hospital / Faculty of Medicine Sebelas Maret University, Surakarta, Indonesia.

<sup>2</sup>Department of Pathological Anatomy, Dr. Moewardi Hospital / Faculty of Medicine Sebelas Maret University, Surakarta, Indonesia.

### Authors' contributions

This work was carried out in collaboration among all authors. Author RDNT designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors AS, KYY, BW, DH and NA managed the analyses of the study. Author IBB managed the literature searches. All authors read and approved the final manuscript.

### Article Information

#### Editor(s):

(1) Prof. Wagih Mommtaz Ghannam, Mansoura University, Egypt.

#### Reviewers:

(1) Yasmin M. Abd El-Aziz, Port Said University, Egypt.

(2) Franco Mutinelli, Istituto Zooprofilattico Sperimentale delle Venezie, Italy.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/67321>

Original Research Article

Received 02 February 2021

Accepted 07 April 2021

Published 09 April 2021

### ABSTRACT

**Background:** The wound healing process is influenced by various substances with ROS having a fundamental role in destroying microorganisms and degrading damaged tissue structures. However, excessive production of ROS can lead to oxidative stress which is the basis of the pathogenesis of chronic, non-healing wounds. Therefore, an antioxidant that can fight oxidative stress is needed by administering propolis extract as an antioxidant to the granulation tissue of male white rats with skin graft model.

**Aims:** This aims study is to determine the effect of propolis extract as an antioxidant and to correlate it by assessing the quality of angiogenesis (MDA and Caspase-3) in granulation tissue in male white rats with skin graft model.

\*Corresponding author: E-mail: [rickydnsturgeon@gmail.com](mailto:rickydnsturgeon@gmail.com);

**Study Design:** Experimental post-test only control.

**Methodology:** This study was an experimental post-test only control group design using 24 male white rats (*Rattus norvegicus*) as experimental animals which were divided into 4 groups consisting of 6 rats in each group. Wounds of granulation tissue were made on the sample by excising the skin on the back with a size of 1x1 cm which was treated for 7 days then and was given skin graft of 2x2 cm from the back of the rat's fat layer. Propolis extract was given at a dose of 200, 400 and 800 mg/kgBW for 7 days after giving the skin graft. MDA serum and caspase-3 expression were examined after giving the propolis extract. The values of malondialdehyde (MDA) and Caspase-3 expression were analyzed by comparative test using 95% confidence interval and  $p < 0.05$  was considered significant.

**Results:** The comparative test of the mean MDA level was  $p < 0.05$  with the post-hoc test showed  $p < 0.05$  between the control group and the 3 doses group of propolis extract. Meanwhile, the comparative test mean of Caspase-3 histoscore showed  $p > 0.05$ , indicating that there was no significant difference between the control group and the 3 doses group of propolis extract.

**Conclusion:** There was a significant difference in the levels of Malondialdehyde (MDA) although there was no significant difference in the expression of Caspase-3 by giving three different doses of propolis extract to the granulation tissue of male rats (*Rattus norvegicus*) with skin graft model.

**Keywords:** Skin graft; white rats; MDA; Caspase-3.

## 1. INTRODUCTION

The prevalence of wounds continues to increase every year. Data from the 2013 Riskesdas reports shows that the prevalence of wounds in Indonesia is 8.2%, with the highest prevalence in South Sulawesi at 12.8% and the lowest in the Jambi area of 4.5% [1]. The high incidence of injuries can become a health problem that requires treatment and a large amount of funds allocated.

Theoretically, the wound will undergo a physiological healing process. The wound healing process occurs through the role of various substances such as ROS, MDA, caspase 9, NF $\kappa$ B, TGF $\beta$  and VEGF. Gorlach et al. [2] stated that Reactive oxygen species (ROS) are associated with the inflammatory process.

ROS represent the principal agents of oxidative stress. Oxidative stress is an imbalance between oxidative and antioxidant events, which causes oxidative reactions; ROS is involved in the production of free radicals, and is a factor responsible for skin aging and disease progression [3]. The fundamental purpose of the formation of high ROS during skin inflammation is to remove and destroy microorganisms that attack and / or degrade the damaged tissue structure. The ROS system is ubiquitous in the aging process, photoaging, inflammation, wound healing, tumorigenesis and other processes in the skin [4].

Excessive production of ROS can lead to oxidative stress, which has been identified as an important feature in the pathogenesis of chronic wounds. Excessive levels of ROS lead to oxidative modification and biomolecular damage, altering lipid / protein / DNA structure and function, inducing irreversible oxidation of reactive thiol protein groups, which are a feature of oxidative stress, and dysregulation of cell signaling pathways, triggering a downstream signaling cascade leading to impaired release of cytokines and exacerbation of skin diseases [5,6].

An antioxidant is needed that will fight oxidative stress so that the wound healing process is not interrupted. Therefore, this study aims to determine the effect of propolis extract as an antioxidant and to correlate it by assessing the quality of angiogenesis in granulation tissue in male white rats with skin graft model.

## 2. MATERIALS AND METHODS

This research is an experimental post-test only control group design with 24 male white rats (*Rattus norvegicus*) as experimental animals. Measurements on test animals were carried out for seven days after giving the treatment. The entire sample was divided into four groups consisting of:

- a. Control group (K)
- b. Group I: Skin Graft model rats given 200 mg/kgBW propolis extract dose

- c. Group II: Skin Graft model rats given 400 mg/kgBW propolis extract dose
- d. Group III: Skin Graft model rats given 800 mg/kgBW propolis extract dose

Each group consisted of 6 male white rats (*Rattus Norvegicus*) treated with Skin Graft and then the levels of malondialdehyde (MDA) and the expression of Caspase-3 granular tissue were measured after treatment.

## 2.1 Provision of Propolis Extract

Propolis was obtained from bee extract with maceration techniques. The bee used was the *Apis mellifera* species. Dry propolis was cleaned and blended until smooth, then weighed as much as 500 grams, and then put into a beaker glass. The maceration process was carried out by adding 3.75 liters of 70% ethanol as a solvent. The mixture of ingredients was stored for seven days in a room that was not exposed to sunlight with strong shaking or stirring with a stirring spatula twice a day [7].

The next step was filtering with a Buchner funnel and filter paper to separate the filtrate from the dregs into the Erlenmeyer flask to obtain the filtrate. The filtration results obtained were evaporated by a rotary evaporator at a temperature of 45°C with vacuum pressure (<1 atm) for approximately 4 hours so that a thick concentration of propolis extract was obtained ( $\pm$  100 g). Furthermore, the propolis extract was evaporated for 24 hours in a beaker glass so that the ethanol content evaporates [8].

Provision of propolis extract in different doses to male white rats (*Rattus norvegicus*) skin graft model was given orally using an oral stomach gavage. The dose of propolis extract used was a dose of 200, 400 mg and 800 mg/kgBW which is given orally every day for 7 days.

## 2.2 Skin Graft Model Mouse Formation

The experimental animal model of granulation tissue wound was formed through the skin excision method (epidermis, dermis and part of the sub-cutis/panniculus carnosus) in the back area with a size of 2 x 2 cm using scalpel No. 10 such as the procedure for taking the skin graft donor with hemostasis using adrenaline 1: 100,000 or using 6-0 monofilament thread ligation. The edges of the wound were anchored with continuous sutures to prevent contraction of the wound edges. The wound was treated using

NaCl 0.9% and left for 7 days without treatment. This model is useful for creating inflammatory wounds of granulation tissue - Extra Cellular Matrix. This model can be induced by either mice or rat and is independent of a particular strain. This model within 2 weeks can cause inflammation in the proliferation phase (granulation tissue) so that inflammatory factors can be investigated.

Skin graft donor was taken in the area around the back with a size of 1 x 1 cm<sup>2</sup>. The skin graft was sutured to the wound with 6-0 monofilament thread and fixed with moist gauze (0.9% NaCl) and cotton for a bolster covered with Tegaderm (3M Company, St. Paul, Minn.). The donor skin graft wound was sutured with 6-0 monofilament suture and the wound was treated with gentamycin skin ointment and covered with Tegaderm (3M Company, St. Paul, Minn.)

## 2.3 MDA Level Examination

Blood MDA levels in this study were serum MDA levels of white rats with skin graft treatment after fasting for 12 hours, which was measured by the Thiobarbituric Acid Reacting Substances (TBARS) method and expressed in units of nmol / mL. Malondialdehyde levels were measured using the KitDiaSys measuring instrument which was done by an expert from the PAU UGM Yogyakarta Animal Laboratory. Blood was drawn from the retro-orbital vein. Measurement of blood MDA levels was carried out after being given the propolis extract, after 7 days of giving the skin graft.

## 2.4 Immunohistochemical Examination of Caspase-3 Expression

Immunohistochemical staining procedures were performed for caspase-3 examination. For immunohistochemical, polyclonal primary antibody, anti-rabbit Ig G, and anti caspase-3 (Thermo Scientific) with 1:100 dilution were used. Antigen was retrieved using citrate buffer with Mayer hematoxylin counterstaining. Caspase-3 expression was calculated by observing cells with no less than 500 cells and more than 5 high power fields. Interpretation criteria are based on the intensity of staining and the percentage of positive cells. The number of cell was determined using the number of cells was determined using the Olympus CX 21 microscope with a small magnification of 40X first followed by medium and large magnification of 100X and 400X respectively. The preparations were then read in sequence from left to right, top to bottom.

## 2.5 Statistical Analysis

The statistical test used was the comparative test by comparing the average levels of MDA and caspase-3 expression using the one-way ANOVA test if the data is normal, and the Kruskal-Wallis comparative test if the data is not normal. All analysis of the research results were analyzed using the statistical application tool IBM-SPSS version 25. P-value <0.05 was considered statistically significant with 95% as the confidence interval value.

## 3. RESULTS AND DISCUSSION

Post-test MDA level data collection was carried out on August 21, 2020 after being given treatment in the form of propolis extract for 7 days. The mean MDA levels after being given skin graft treatment are shown in Table 1.

The results in Table 1 show that there are differences in MDA levels given skin graft treatment when compared to MDA levels given skin grafts with propolis extract. The MDA level with skin graft treatment was 8.050 nmol/ml while the mean MDA level with skin graft treatment and 200 mg/kgBW of propolis extract showed a difference of 6.282 nmol/ml. Giving skin graft and

propolis extract treatment with a larger dose of 400 mg/kgBW made the average MDA level to 2.407 nmol/ml, and giving the propolis extract with the largest dose, 800 mg/kgBW, showed a much greater difference in MDA levels, compared to the control group given 1.425 nmol/ml. From Table 1, it is obtained a p-value of < 0.05, which indicates that at least two groups have significantly different mean MDA levels. Post-hoc test analysis was conducted to find out which groups had differences. The results of the post-hoc test can be seen in Table 2.

From the results of the post-hoc analysis, it was found that there were significant differences in the mean levels of MDA between group K and groups P1, P2 and P3 (p-value < 0.05) and between groups P1 with groups P2 and P3. These results indicate that the differences in the results of existing MDA levels were significant between group K and groups P1, P2, and P3 and group P1 with groups P2 and P3. On the other hand, there was no significant difference in the results of MDA levels between group P2 and group P3 (p > 0.05). These results indicate that the difference in the results of MDA levels between groups P2 and P3 was not statistically significant.

**Table 1. Results of the analysis of MDA levels using the one-way ANOVA test**

Group	N	Mean ± Standard deviation (nmol/ml)	P-value
K	6	8.050 ± 0.536	0.000
P1	6	6.282 ± 0.192	
P2	6	2.407 ± 0.792	
P3	6	1.452 ± 0.153	

**Table 2. Results of analysis of MDA levels using the one-way ANOVA test**

Group	P-value
MDA levels	
Group K vs Group P1	0.001
Group K vs Group P2	0.000
Group K vs Group P3	0.000
Group P1 vs Group P2	0.000
Group P1 vs Group P3	0.000
Group P2 vs Group P3	0.157

\*Group K: Skin graft control

Group P1: Skin graft with 200 mg /kgBW propolis extract

Group P2: Skin graft with 400 mg/kgBW of propolis extract

Group P3: Skin graft with 800 mg/kgBW of propolis extract

**Table 3. Results of Caspase-3 histoscore analysis using the Kruskal-Wallis test**

Group	N	Mean ± Standard deviation	P value
K	6	10.00 ± 2,19	0.557
P1	6	9.83 ± 3,71	
P2	6	8.50 ± 3,33	
P3	6	8.67 ± 1,63	

The results in Table 3 show that there is a difference between Caspase-3 histoscore given skin graft treatment and Caspase-3 histoscore given skin graft treatment with propolis extract. The mean of Caspase-3 histoscore treated with skin graft was 10.0 while the mean Caspase-3 histoscore treated with skin graft and 200 mg/kgBW of propolis extract had a mean of 9.83. Giving skin graft and propolis extract treatment with a larger dose of 400 mg/kgBW made the average Caspase-3 histoscore to be 8.50. Finally, the administration of propolis extract with the largest dose, 800 mg/kgBW, showed a significant difference of Caspase-3 histoscore compared to the control group.

The distribution of Caspase-3 histoscore data was checked using the Shapiro-Wilk test. The results of the normality test showed that the data were not normally distributed. Thus, the comparative test of mean histoscore score in each trial group was carried out using the Kruskal-Wallis test. The results of the Kruskal-Wallis mean comparative test can be seen in Table 3.

Based on this study, it was found that there were differences between the MDA levels in the control group (K) and the group given 200, 400 and 800 mg/kgBW of propolis extract. The results also showed that the administration of 400 mg/kgBW of propolis extract gave the most statistically significant difference in MDA levels because there was no statistically significant difference between the administrations of 400 and 800 mg/kgBW of propolis extract.

In a study conducted by Damayanti et al. [9] which compared MDA levels in the administration of propolis doses to the brain of a rat brain injury model due to trauma, it was found that there was a difference between the control group rat and the rat given propolis although there was no significant difference between the doses of 50, 100 and 200 mg/kgBW. The researcher explained that the higher dose of propolis extract was more likely to reduce MDA levels in the brain tissue of rats in the traumatic brain injury model, but the insignificant difference was probably due to the too low dose range of propolis administration. Meanwhile, based on the correlation test, it was found that there was a weak negative correlation between the doses, indicating that the higher dose of propolis extract may decrease MDA levels. The results obtained by this study indicate that between the dosages of propolis extract of 50, 100, and 200 mg/kgBW,

there is no significant difference in MDA levels, but Damayanti et al. suggest using even higher doses of MDA for future research [9].

The results obtained are in line with another study regarding the effect of reducing MDA levels by propolis extract conducted by Hoesada et al. [10] in their research on propolis extract with a total of 14.5, 27, and 41.5 mg/kgBW against MDA levels in male Wistar rats treated with cigarette smoke for 21 days. The results obtained indicate that the propolis extract can significantly reduce MDA levels in male Wistar rats treated with cigarette smoke [10]. In addition, in a study conducted by Prasetyo et al. [11] which examined the effects of ethanol extract of propolis on the levels of inflammation and serum MDA in sepsis model mice, MDA levels of sepsis model mice treated with propolis extract decreased drastically from 0.39 to 0.09 micromole/L. The results showed that the ethanol extract of propolis could reduce serum MDA levels in sepsis model experimental animals [11].

The effect of this difference in MDA levels occurs due to the presence of antioxidants from propolis such as terpenoids, flavonoids and cinnamic acid. The antioxidant property of flavonoids is that they can take free radicals (ROS) directly by donating hydrogen atoms or transferring electrons. In addition, flavonoids can also act as intracellular antioxidants through inhibition of free radical-forming enzymes such as xanthine oxidase (XO), lipoxygenase, protein kinase C, cyclooxygenase (COX), monooxygenase, and NADPH oxidase [12]. This is also in line with the research conducted by Hardiningtyas et al. [13] examining the antioxidant activity of the white apiary mangrove leaves and finding the phytochemical content of ethyl acetate extract from the leaves of the white fire mangrove which was rich in flavonoids, steroids, and triterpenoids. The extract was then given to Sprague Dawley rats which were hepatotoxic induced with carbon tetrachloride CCL4 which caused high levels of liver MDA and then administered with ethyl acetate extract of white fire mangrove leaves. There was a statistically significant difference indicating a protective effect and inhibition of lipid peroxidation from Ethyl acetate extract of the white flame mangrove leaves which is rich in these flavonoids [13].

Based on the measurement of Caspase-3 levels after 7 days, the experimental animals were given skin graft and propolis extract treatment. The Kruskal-Wallis mean comparative analysis

test showed  $p > 0.05$ . This shows that no two groups have different Caspase-3 means which are statistically significant [14].

The Caspase-3 protein has an important influence on apoptosis. Caspase-3 is expressed in normal tissue and significantly lowers in gastric cancer. Another study also showed that Caspase-3 expression decreased in tongue cancer with overexpression of survivin which had an effect on inhibition of caspase-3 expression [15]. In the process of angiogenesis, VEGF is expressed and increases the expression of survivin which can inhibit apoptosis through the apoptotic signaling pathway, thereby protecting the newly arising immature tumor vascular endothelial cells, enabling them to avoid apoptosis and subsequently producing more VEGF [16].

The results obtained in this study are not in line with the research conducted by Tandean et al. [17] on Sprague Dawley rats that were treated with closed head injury and then given 200 mg/kgBW of propolis extract for 7 days, showing that the administration of propolis extract reduced caspase-3 expression which was statistically significant through the caspase-dependent pathway and caspase-independent pathway inhibition mechanisms. The results obtained are also different from another study conducted by Sun et al. [18] which examined the effects of propolis extract on a Neuro-2a neuroblastoma (N2a) cell model that was given hypoxia and hypoglycemia treatment, showing that propolis extract could downregulate Caspase-3.

The results of the comparative test between the control group and the group given propolis extract, which was not statistically significant, still showed a decrease in Caspase-3 expression in the latter group. This is due to the contents of the propolis extract, namely Chrysin, pinocembrin, pinobanksin, apigenin, and galangin which apparently show an inhibitory effect on the apoptotic pathway such as reducing the expression of Caspase-3 and Bax and increasing the expression of Bcl-2 mRNA [18].

#### 4. CONCLUSION

There was a significant difference in malondialdehyde (MDA) levels in the administration of three different doses of propolis extract, and there was an insignificant difference in the expression of Caspase-3 in giving propolis

extract to the granulation tissue of the skin graft model male rats. The most statistically significant difference in MDA levels occurred when the sample was given propolis extract at a dose of 400 mg/kgBW.

#### CONSENT

It is not applicable.

#### ETHICAL APPROVAL

This research obtained ethical approval from the Research Ethics Committee of RSUD Dr. Moewardi Surakarta with number 721/VI/HREC/2020. All authors hereby declare that "Principles of laboratory animal care" (NIH publication No. 85-23, revised 1985) were followed, as well as specific national laws where applicable. All experiments have been examined and approved by the appropriate ethics committee.

#### ACKNOWLEDGEMENTS

The author's gratitude to the Department of Anatomical Pathology Faculty of Medicine Sebelas Maret University and Department of Surgery Dr. Moewardi Hospital Surakarta who have supported this research. The author is responsible for funding the study without the involvement of grants or other sources of funding.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Riskesdas. Riset Kesehatan Dasar. Departemen Kesehatan Indonesia; 2013. Bahasa
2. Görlach A, Dimova EY, Petry A, Martínez-Ruiz A, Hernansanz-Agustín P, Rolo AP, et al. Reactive oxygen species, nutrition, hypoxia and diseases: Problems solved? *Redox Biology*. Elsevier B.V. 2015;6:372–85. Accessed 8 August 2020 Available: <https://pubmed.ncbi.nlm.nih.gov/26339717/>
3. Bickers DR, Athar M. Oxidative stress in the pathogenesis of skin disease. *J Invest Dermatol*. 2006;126(12):2565–75.

4. Masaki H. Role of antioxidants in the skin: Anti-aging effects. *Journal of Dermatological Science*. 2010;58:85-90. Accessed 9 August 2020 Available: <https://pubmed.ncbi.nlm.nih.gov/20399614/>
5. Moseley R, Hilton JR, Waddington RJ, Harding KG, Stephens P, Thomas DW. Comparison of oxidative stress biomarker profiles between acute and chronic wound environments. *Wound Repair Regen*. 2004;12(4):419–29. Accessed 9 August 2020 Available: <https://pubmed.ncbi.nlm.nih.gov/15260807/>
6. Xu H, Zheng Y-W, Liu Q, Liu L-P, Luo F-L, Zhou H-C, et al. Reactive oxygen species in skin repair, regeneration, aging and inflammation. In: *Reactive Oxygen Species (ROS) in Living Cell*. InTech; 2018. Accessed 9 August 2020 Available: <http://dx.doi.org/10.5772/intechopen.72747>
7. Bankova V, Trusheva B, Popova M. New developments in propolis chemical diversity studies (Since 2000); 2008.
8. Sarsono, Syarifah I, Martini, Prasetyo DH. Identifikasi caffeic acid phenethyl ester dalam ekstrak etanol propolis isolat gunung lawu. *JBAI*. 2012;8(2):132. Bahasa
9. Damayanti R, Enggar Fitri L, Dalhar M. Pengaruh Pemberian Propolis terhadap Ekspresi INOS dan Kadar MDA pada Otak Tikus Model Cedera Otak Traumatik. *J Kedokt Brawijaya*. 2016;29(2):110–6. Bahasa
10. Hoesada I, Nasihun T, Isradji I. The effect of propolis extract on MDA levels (Malondialdehyde) and sperm quality on epididimis (Experimental Study on Wistar Strain Male Rats Exposed to Kretek Cigarettes). *Sains Med J Kedokt dan Kesehat*. 2016;7(1):09-14. Accessed 6 September 2020 Available: <http://jurnal.unissula.ac.id/index.php/sainsmedika/article/view/1165>
11. Prasetyo DH, Suparyanti EL, Guntur A. Ekstrak Etanol Propolis Isolat Menurunkan Derajat Inflamasi dan Kadar Malondialdehid pada Serum Tikus Model Sepsis Ethanol extract of Propolis Reduces the Level of Inflammation and Serum Malondialdehyde in Sepsis Rats Model. *Mkb*. 2013;45(3):161–6. Bahasa
12. Gordon MH, Roedig-Penman A. Antioxidant properties of flavonols. *Lipids Heal Nutr*. 2010;23(4):47–64.
13. Hardiningtyas SD, Purwaningsih S, Handharyani E. Aktivitas Antioksidan Dan Efek Hepatoprotektif Daun Bakau Api-Api Putih. *J Pengolah Has Perikan Indones*. 2014;17(1):80–91. Bahasa
14. Dahlan S. Langkah-langkah membuat proposal penelitian bidang kedokteran dan menerapkan hasil penelitian; 2011. Bahasa
15. Xu JH, Wang AX, Huang HZ, Wang JG, Pan C. Bin, Zhang B. Survivin shRNA induces caspase-3-dependent apoptosis and enhances cisplatin sensitivity in squamous cell carcinoma of the tongue. *Oncol Res*. 2010;18(8):377–85. Accessed 25 November 2020 Available: <https://pubmed.ncbi.nlm.nih.gov/20441052/>
16. Wang S, Xu J, Zhang Q. Clinical significance of survivin and vascular endothelial growth factor mRNA detection in the peripheral whole blood of breast cancer patients. *Neoplasma*. 2016;63(1):133–40. Accessed 25 November 2020 Available: <https://pubmed.ncbi.nlm.nih.gov/26639243/>
17. Tandean S, Japardi I, Loe ML, Riawan W, July J. Protective effects of propolis extract in a rat model of traumatic brain injury via hsp70 induction. *Open Access Maced J Med Sci*. 2019;7(17):2763–6.
18. Sun LP, Xu X, Hwang HH, Wang X, Su KY, Chen YLS. Dichloromethane extracts of propolis protect cell from oxygen glucose deprivation-induced oxidative stress via reducing apoptosis. *Food Nutr Res*. 2016;60. Accessed 7 September 2020 Available: <http://dx.doi.org/10.3402/fnr.v60.30081>

© 2021 Tyastono et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
The peer review history for this paper can be accessed here:  
<http://www.sdiarticle4.com/review-history/67321>