



Comparison of Non-linear Growth Curve Models in Non-descript California and New Zealand Rabbits Reared in the Tropical Conditions of Nigeria

Okeke Rufina Obioma¹, Suleiman Ibrahim Onotu², Omotugba Stephen Kayode³,
Ibikunle Kehinde Yemiola⁴, Idris Abdullahi⁵, Louis Ugwu⁶,
Adeniyi Adebayo Kunle⁷ and Oludayo Michael Akinsola^{8*}

¹Department of Animal Science, Ahmadu Bello University, Nigeria.

²Department of Animal Science, Faculty of Agriculture, Bayero University, Kano, Nigeria.

³Department of Basic Science, Federal College of Wildlife Management, Nigeria.

⁴National Institute for Hospitality and Tourism, Edo State, Nigeria.

⁵National Animal Production Research Institute, Ahmadu Bello University, Zaria, Nigeria.

⁶Department of Biology, School of Science, Federal College of Education, Katsina, Nigeria.

⁷Department of Animal Production, Faculty of Agriculture, University of Jos, Nigeria.

⁸Department of Theriogenology and Production, Faculty of Veterinary Medicine, University of Jos, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. Author ORO designed the study. Author SIO performed the statistical analysis. Author OSK wrote the protocol. Author IKY wrote the first draft of the manuscript. Authors IA and OMA managed the analyses of the study. Author AAK managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJRAF/2020/v5i230080

Editor(s):

(1) Dr. Cengiz Yucedag, Professor, Department of Landscape Architecture, Mehmet Akif Ersoy University, Turkey.

Reviewers:

(1) Ahmed N. F. Neamat-Allah, Zagazig University, Egypt.

(2) Patricia Ferreira Ponciano Ferraz, Federal University of Lavras, Brazil.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/54482>

Original Research Article

Received 07 December 2019

Accepted 12 February 2020

Published 28 February 2020

ABSTRACT

Nonlinear functions of body weight at different age intervals were used to estimate the growth pattern in New Zealand White and California rabbits. Gompertz and Logistic functions of 3 and 4 parameters were fitted to Age-weight data matrix. Age-weight records of New Zealand White and California rabbits from birth were monitored to 56 days to estimate the average growth curve for

*Corresponding author: E-mail: dayoakinsola@gmail.com;

each breed. The weight difference between breeds was consistently in favor of California rabbits as compared to New Zealand White. It was concluded that the Gompertz and logistic models were both parsimonious and adequate in describing the growth patterns of New Zealand White and California rabbits in the tropical conditions of Nigeria.

Keywords: Gompertz model; logistic models; non-linear growth curve models; rabbits; nonlinear functions; body weight.

1. INTRODUCTION

The medium rabbits in the tropical countries serve as a good source of cheap animal protein due to quest for protein sufficiency among the teeming populace of Sub Saharan African [1]. Rabbits possess unique biological characteristics such as high biological value amino acids, meat low in calories and fat, prolificacy, short gestation length, initial capital outlay, great genetic flexibility, highly unsaturated lipids (60% of the total fatty acids), valuable animal model for biomedical research, low sodium and cholesterol levels [1]. Weight changes data are used by farmers to monitor on the spot assessment of health status of rabbit for forecasting growth trait within trajectory of time. Growth is a complex trait of interest in domestic animals and is as expressed as the increase in size, number, or mass with time. Growth could be expressed as the increase in size number or weight changes.

The use of maximum-likelihood solving estimates in reducing the model noise using discrete and continuous data in model fitting have been reported by several studies. The use of Gompertz and logistic models were used to analyse the growth of rabbit [2], lamb [3], pig [4] and cattle [5,6]. Several researches had modeled rabbit growth curve using logarithmic model [7], stochastic model [8] and a general linear mixed model [9].

Some selected mathematical functions commonly used in describing the growth phase of rabbits include the Gompertz, Logistic, Brody, von Bertalanffy and Richards growth models [9]. The growth functions can be grouped into three main categories: those with a diminishing returns behaviour (Brody model), those with sigmoidal shape and a fixed inflexion point (Gompertz, Logistic and von Bertalanffy models) and those with a flexible inflexion point (Richards model). The Logistic, Gompertz and von Bertalanffy models exhibit inflection points at about 50, 37 and 30% of the mature weight (asymptote), respectively.

Therefore, the objective of this study was to define the most appropriate nonlinear model to estimate California and New Zealand rabbit breeds bodyweight changes over time trajectory in Northern Guinea Savannah Zone of Nigeria.

2. MATERIALS AND METHODS

The forty animals (n=20 California and n=20 New Zealand White) used in this experiment originated from the Dagwom experimental rabbit farm at National Veterinary Research Institute, Vom, Jos, Plateau State. The rabbits were kept in individual cages and were fed with conventional formulated growers feed and grasses such as *Calopogonium mucunoides* and *Panicum maximum ad libitum*. Fresh drinking water was also provided always to the rabbits. Other daily routine management practices were carried out regularly. Bodyweight of rabbits were taken by digital weighing scale (Mettler Toledo, Top Pan Sensitive Balance, J. Liang Int. Ltd. U.K.) on weekly basis.

2.1 Statistical Procedures

Bodyweights at different ages were fitted to 3-parameters and 4-parameters models. The three-parameter functions sought to find a way to fit a response that is bound between zero and the estimated asymptote. The four-parameter functions are bounded between two estimated asymptotes. Age and Inflection point were computed as described in the non linear algorithm of JMP 13 statistical software. The following growth curve models were used to determine the parameters belonging to the growth curves (Table 1).

The model performance was compared based on the coefficient of determination (R^2), root means square error (RMSE), Akaike information criterion (AIC) and Bayesian information criterion (BIC). RMSE and R^2 of models were considered for the calculated goodness of fit parameters for the comparison among the different models between breeds of rabbits. Coefficient of determination (R^2) = 1-(SSE/SST) where SSE is the sum of square of errors, and SST is the total

Table 1. Functional form of growth curve models

Models	Functions
Logistic 3P ¹	$BW = a \cdot \text{Exp} \left(-\text{Exp}(-b \cdot (\text{Age} - c)) \right)$
Gompertz 3P ²	$BW = \frac{c}{\left(1 + \text{Exp}(-a \cdot (\text{Age} - b)) \right)}$
Logistic 4P ³	$BW = a + (b - a) \cdot \text{Exp} \left(-\text{Exp}(-c \cdot (\text{Age} - d)) \right)$
Gompertz 4P ⁴	$BW = \frac{(d - c)}{\left(1 + \text{Exp}(-a \cdot (\text{Age} - b)) \right)}$

P=parameter; a=growth rate; b=inflexion point; c=lower asymptote; d=upper asymptote; BW=Bodyweight; ¹[4]; ²[10]; ³[11]; ⁴[12]

sum of squares. Akaike's information criteria (AIC) = n*ln (SSE/n) + 2k, where n; the number of observations, SSE; sum of square of errors, k; the number of parameters.

Schwarz Bayesian information criterion (BIC) = n*ln(SSE/n) + k*ln(n), where n; the number of observations, SSE; sum of square of errors, k; the number of parameters [13]. The growth data was fitted using the prediction profiler interface of JMP 13.2 software. Estimated mean parameters obtained from different models were used to plot general growth curves of the non-descript rabbits.

3. RESULTS

Table 2 shows the selection criteria of differently parameterized mathematical models in non-descript New Zealand White and California Rabbits. All the models were parsimonious with reduced model noise in reconstructing the growth curve between the breeds. Logistic 4P showed the highest coefficient of determination of R²=0.989 and 0.992 in New Zealand and

California Rabbits. Gompertz 3P and Gompertz 4P had similar prediction efficiencies in the New Zealand White (R²=0.988) while Gompertz and Logistic 4P showed similar values in California Rabbits (R²=0.992). The best Bayesian information criterion and Akaike information criterion was recorded in Gompertz 3P (0.56 and 111.04) for New Zealand White and California rabbits (98.20 and 107.41). Gompertz 3P had most reduced best model noise in New Zealand White while Logistic 4P recorded the least noise in California Rabbit.

The growth curves parameter estimate in Non-descript New Zealand White and California Rabbits are presented in Table 3 and Figs. 1 and 2. The optimal age for maturity was at 28 days with bodyweight ranging from 1006.94 – 3522.14 g in New Zealand White and 2544.88 – 3639.49 g in California rabbits. The Gompertz 3P and 4P showed the best age at inflection point in New Zealand White (13.40 days) and California rabbits (28.84 days). The California rabbits showed early maturing/growth rate (0.04 – 0.06) than New Zealand White (0.01 – 0.06).

Table 2. Selection criteria (Weight in grams) for differently parameterized mathematical models in non-descript New Zealand and California White Rabbits

Model/Breed	AIC	BIC	SSE	MSE	RMSE	R ²
New Zealand white						
Gompertz 3P	111.04	101.82	16272.44	2712.07	52.07	0.988
Logistic 3P	111.57	102.36	17271.94	2878.65	53.65	0.987
Logistic 4P	121.96	102.95	14446.30	2889.26	53.75	0.989
Gompertz 4P	122.56	103.55	15434.23	3086.84	55.55	0.988
California						
Gompertz 3P	107.41	98.20	10880.82	1813.46	42.58	0.990
Logistic 3P	108.58	99.37	12387.85	2064.64	45.43	0.989
Logistic 4P	117.62	98.61	8916.57	1783.31	42.22	0.992
Gompertz 4P	117.95	98.94	9247.95	1849.59	43.00	0.992

P=parameters; AIC=Akaike Information Criterion, BIC= Bayesian Information Criterion, SSE-Sum of square error, MSE-Mean of square error, RMSE-Root mean square error, R²=Coefficient of Determination




Fig. 1. Curve fitting for body weight changes in relations with age in New Zealand white rabbit




Fig. 2. Curve fitting for body weight changes in relations with age in California rabbit

Table 3. Growth curves parameter estimate in non-descript New Zealand and California white rabbits

Models/Breed	Growth rate	Inflection point	Optimal age (weeks)	First derivative	Second derivative	Optimal weight (grams)
New Zealand						
Gompertz 3P	0.01	13.40	28	18.10	0.68	1795.17
Logistic 3P	0.06	41.38	28	17.78	0.76	2776.18
Logistic 4P	0.03	33.01	28	17.70	0.56	1006.94
Gompertz 4P	0.04	25.25	28	17.96	0.59	3522.14
California						
Gompertz 3P	0.06	39.86	28	16.55	0.85	3629.49
Logistic 3P	0.06	31.30	28	16.09	0.87	2980.89
Logistic 4P	0.04	30.88	28	16.00	0.82	2650.22
Gompertz 4P	0.04	28.84	28	15.89	0.80	2544.88

P=Parameter

4. DISCUSSION

Genetic gain is directly related to economics of production which have attracted the attention of geneticist, livestock scientists and private rabbit enterprises. The utilization of these parameters in growth models using age-weight gain matrix data could influence economic returns positively. The higher coefficients of determination recorded for both breeds were comparable with observations of several researchers about the growth of rabbits and these were analysed with Logarithmic model [7] Stochastic model [8] and General Linear Mixed Model [9]. The higher maturity values observed in California rabbits research about the growth of rabbits were analyzed agrees with the report of Kurnianto et al. [14] who showed that large values of k indicated early maturing individuals, whereas small k values indicated late maturing individuals. Point of inflection was correspondent to two parameters, namely age at point inflection (t_i) and weight at point of inflection (y_i). The rabbit breed with higher *growth rate* value showed higher point of inflection. Differences in growth patterns were expressed in terms of differences in mature weight and maturing rate in both New Zealand White and California rabbits. The optimal weight in New Zealand White rabbit showed larger fluctuations as compared to California rabbits. The California rabbit was heavier and also matured earlier than the New Zealand White rabbits. Comparisons of asymptotic body weight obtained with different functions showed that the Gompertz 4P in New Zealand White and Gompertz 3P model in California rabbit provided better weight estimates. Akinsola [1] reported variations in prediction model estimates in different genotypes

of rabbit. The Logistic and Gompertz models are more accurate in predicting the body weight of these animals and have better overall fit. The prediction of adult BW using nonlinear functions can be accurate when growth curve parameters and their covariance components are estimated jointly [6]. The estimation of potential final weight in different species is a function of an algorithm fitting and the accuracy of judging is possible only when precise final weight is available [3].

5. CONCLUSION

The Gompertz and Logistic models were both parsimonious and adequate for describing the growth patterns of New Zealand White and California rabbits. The California rabbit were heavier and mature than New Zealand White rabbit during the growth phase. In comparison, the time taken to reach maturity was longer for New Zealand White rabbit as compared to California rabbits.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Akinsola OM. Genetic and physiological evaluation of Hyla rabbits in Guinea Savannah Zone of Nigeria. Zaria: M.Sc. Dissertation Submitted to the Department of Animal Science, Ahmadu Bello University. 2012;66–70.
2. Blasco A, Piles M, Varona L. A Bayesian analysis of the effect of selection for growth rate on growth curves in rabbits. Genet. Sel. Evol. 2003;35:21-41.

3. Lambe NR, Navajas EA, Simm G, Bungler L. A genetic investigation of various growth models to describe growth of lambs of two contrasting breeds. *J. Anim. Sci.* 2006; 84:2642-2654.
4. Strathe AB, Danfear A, Sorensen H, Kebreab E. A multilevel nonlinear mixed-effects approach to model growth in pigs. *J. Anim. Sci.* 2010;88:638-649.
5. Budimulyati LS, Noor RR, Saefuddin A, Talib C. Comparison on accuracy of Logistic, Gompertz and Von Bertalanffy models in predicting growth of new born calf until first mating of Holstein Friesian heifers. *J. Indonesian Trop. Anim. Agric.* 2012;37(3):151-160.
6. Forni S, Piles M, Blasco A, Varona L, Oliveir HN, Lobo RB. Comparison of different nonlinear functions to describe Nelore cattle growth. *J. Anim. Sci.* 2009; 87:496-506.
7. Rao DR, Sunki GR, Johnson WM, Chen CP. Postnatal growth of New Zealand White rabbit (*Oryctolagus cuniculus*). *J. Anim. Sci.* 1997;44:1021-1025.
8. Sampaio IMB, Ferreira WM, Bastos AF. The use of a stochastic model of rabbit growth for culling. *World Rabbit Sci.* 2005; 13:107-112.
9. McNitt JI, Lukehfar SD. Breed and environmental effects on postweaning growth of rabbits. *J. Anim. Sci.* 2005;71: 1996-2005.
10. Gompertz B. On the nature of the function expressive of the law of human mortality, and on a new mode of determining the value of life contingencies. *Phil. Trans. Roy. Soc. London.* 1832;123:513-585.
11. Ratkowsky DA, Reedy TJ. Choosing near-linear parameters in the four-parameter logistic model for radioligand and related assays. *Biometrics.* 1986;42:575-582.
12. Gibson AM, Bratchell N, Roberts TA. The effect of sodium chloride and temperature on the rate and extent of growth of *Clostridium botulinum* type A in pasteurized pork slurry. *Journal of Applied Bacteriology.* 1987;62(6):479-90. [PMID: 3305458]
13. Burnham Kenneth P, David R. Anderson. Model selection and multimodel inference: A practical information-theoretical approach. 2nd Ed. New York: Springer Verlag; 2002.
14. Kurnianto E, Shinjo A, Suga D. Analysis of growth in intersubspecific crossing of mice using Gompertz model. *Asian-Aust. J. Anim. Sci.* 1998;11:84-88.

© 2020 Obioma et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/54482>