

Acute left main coronary artery occlusion

K. Mehmet Burgazli¹, Mehmet Bilgin²,
Nedim Soydan³, Ridvan Chasan⁴, Ali Erdogan⁵

ABSTRACT

The treatment of an acute left main coronary artery occlusion still poses a challenge. In this case report we present a 50-year-old patient with an acute occlusion of the left main artery. After a successful angioplasty without “stenting” due to the complexity of the stenosis the patient underwent a successful bypass surgery. We discuss the therapeutic options of acute left main occlusion regarding medical, interventional and surgical options.

KEY WORDS: Acute left main occlusion, Percutaneous coronary angioplasty, Bypass surgery.

doi: <http://dx.doi.org/10.12669/pjms.291.2819>

How to cite this:

Burgazli KM, Bilgin M, Soydan N, Chasan R, Erdogan A. Acute left main coronary artery occlusion. *Pak J Med Sci* 2013;29(1):216-217.
doi: <http://dx.doi.org/10.12669/pjms.291.2819>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Acute left main coronary artery occlusion presents a high-risk situation. The medical treatment of the unprotected left main coronary artery has high mortality.¹ Besides the bypass surgery with an evident benefit in comparison with medical therapy, the percutaneous transluminal coronary angioplasty (PTCA) offers another therapeutic option.²

CASE REPORT

A 50-year old male patient was admitted to our hospital in cardiogenic shock with ST-elevation myocardial infarction (STEMI). The cardiovascular risk factors were dyslipidemia, hypertension, and smoking. The coronary angiography (Fig. 1-3) presented a two vessel disease with a high-grade left main coronary artery occlusion {100% de-novo stenosis in the left main}. After successful

1. K. Mehmet Burgazli,
Wuppertal Research and Medical Center,
Department of Innere Medizin,
Angiology Wuppertal, Germany.
2. Mehmet Bilgin,
Department of Radiology,
Bezmialem Vakif University, Istanbul, Turkey.
3. Nedim Soydan,
4. Ridvan Chasan ,
5. Ali Erdogan,
- 1,3-5: University Clinic of Giessen,
Internal Medicine,
Cardiology, Angiology, Giessen,
Germany.

Correspondence:

Dr. K. Mehmet Burgazli,
Medical and Research Center Wuppertal,
Department of Internal Medicine, A
ngiology, Phlebology,
Wichlinghauser Str. 110/11242277,
Wuppertal, Germany.
E-mail: mehmetburgazli@hotmail.de

- * Received for Publication: August 15, 2012
- * Revision Received: October 16, 2012
- * Revision Accepted: October 16, 2012




Fig.1: Occlusion of the left main.




Fig.2: After single balloon dilatation of the left main.

PTCA the stenosis was reduced to 70%. The patient was supported with an intra-aortic balloon counterpulsation (IABP) device during phase of cardiac decompensation with acute pulmonary edema. He received low dose of catecholamines. The echocardiography documented a severely reduced left ventricular ejection fraction. After stabilization in our intensive care unit, the patient was then transferred to cardiac surgery. After coronary artery bypass grafting (CABG; left Arteria mammaria to LAD, Vena saphena magna to Ramus intermedius and to Ramus circumflexus) patient died the 11th postoperative day due to septic shock.

DISCUSSION

Acute occlusion of the left main coronary artery is rare and generally fatal. The mechanism is mostly acute thrombosis and the clinical presentation shows an extensive infarction usually with cardiogenic shock. Bypass surgery shows a survival benefit compared to medical therapy, and is the current standard of therapy for patients with unprotected left main coronary artery occlusion.³ The initial attempt of the percutaneous coronary intervention (PCI) documented frequent serious comorbidities and high event rates. In contrast the additional stenting of the unprotected left main coronary artery results in better prognosis. Since the appearance of drug eluting stents (DES) with reduction in restenosis rates, recent studies compared PCI with DES versus bypass surgery. In this connection PCI shows non-significant results with lower 6-month⁴ and 12-month mortality and myocardial infarction rates.⁵

Nevertheless, there also exists data with significantly higher 6-month⁴ and 12-month target




Fig.3: Right coronary artery.

vessel revascularization rates,⁵ due to instant restenosis. To our knowledge, our case report represents one of the first time a patient with an acute left main coronary artery occlusion within STEMI, who received a successful PCI for the myocardial revascularization. In this case the PCI is used as a bridge over to CABG. The fact that the patient died the 11th postoperative day due to non cardiac death, signifies basically the acute resolution. Patients with renal dysfunction, prior bypass surgery, advanced age, and severe heart failure are at highest risk. As mentioned our patient presented a highly reduced left ventricular ejection fraction. We may advise PCI as an alternative for patients who are ineligible for bypass surgery, or for emergency myocardial infarction as a bridge over to CABG in selected patients with acute left main occlusion like in our case report.

REFERENCES

1. Cohen MV, Gorlin R. Main left coronary artery disease. Clinical experience from 1964-1974. *Circulation*. 1975;52(2):275-285.
2. Grüntzig AR. Transluminal dilatation of coronary-artery stenosis. *Lancet*. 1978;1(8053):263.
3. Smith SC Jr, Feldman TE, Hirshfeld JW Jr, Jacobs AK, Kern MJ, King SB, et al. ACC/AHA/SCAI 2005 guideline update for percutaneous coronary intervention: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (ACC/AHA/SCAI Writing Committee to Update 2001 Guidelines for Percutaneous Coronary Intervention. *Circulation*. 2006;113(7):e166-e286.
4. Lee MS, Kapoor N, Jamal F, Czer L, Aragon J, Forrester J, et al. Comparison of coronary artery bypass surgery with percutaneous coronary intervention with drug-eluting stents for unprotected left main coronary artery disease. *J Am Coll Cardiol*. 2006;47(4):864-870.
5. Palmerini T, Marzocchi A, Marzocchi C, Ortolani P, Saia F, Savini C, et al. Comparison between coronary angioplasty & coronary artery bypass surgery for the treatment of unprotected left main coronary artery stenosis (the Bologna Registry). *Am J Cardiol*. 2006;98(1):54-59.