

Journal of Global Ecology and Environment

Volume 20, Issue 3, Page 55-67, 2024; Article no.JOGEE.12348 ISSN: 2454-2644

Assessing the Impact of Water Quality and Its Significance on Women's Health in Guatemala

Chelsea Lim ^{a*}

^a Environmental Sciences Division/STEM Science Center, 111 Charlotte Place/Englewood Cliffs, NJ 07632, USA.

Author's contribution

The sole author designed, analysed, interpreted and prepared the manuscript.

Article Information

DOI: https://doi.org/10.56557/jogee/2024/v20i38849

Open Peer Review History: This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: https://prh.ikprress.org/review-history/12348

Original Research Article

Received: 27/06/2024 Accepted: 01/09/2024 Published: 05/09/2024

ABSTRACT

The lack of good quality water in many third-world countries such as Guatemala is a pressing issue that demands immediate attention, among other environmental issues. Although the dangerous effects of poor water quality on children's health are well known and extensively studied, there is relatively less research focusing on water quality's impact on women's health and living conditions. Women in Guatemala are potentially more vulnerable to exposure to toxins and adverse environmental conditions that affect their health differently from men due to diet, behavior, and living conditions. These toxins exposure might lead to various health problems, including urinary tract infections (UTIs), dysmenorrhea, irritable bowel syndrome (IBS), parasites, and eczema. This paper investigates the current state of water quality in Guatemala through a comprehensive literature review of previous studies. The lifestyle factors affecting women's health, including age,

housing, childbirth, health conditions, and access to medical care, were examined using a set of

questionnaire research. Additionally, the survey data from local women in Guatemala provide

*Corresponding author: E-mail: Clim@STEMsc.org;

Cite as: Lim, Chelsea. 2024. "Assessing the Impact of Water Quality and Its Significance on Women's Health in Guatemala". Journal of Global Ecology and Environment 20 (3):55-67. https://doi.org/10.56557/jogee/2024/v20i38849.

practical insights into their living conditions and equip us with a deeper understanding of their challenges. The survey findings suggest that over half of the women rate their overall health fairly and face significant challenges in obtaining medical care.

Keywords: Environmental impact; third world countries; water quality in Guatemala; women's health.

1. INTRODUCTION

Clean water is one of the basic elements for survival and existence in any growing area globally [1,2]. It's a fundamental prerequisite for drinking, hygiene, sanitation, cultivation, and daily life in every aspect [3,4]. Without clean water, communities face challenges that hinder progress and well-being [5]. About 85% of illnesses in developing countries are linked to poor water and sanitation conditions [6]. 1 out of every 5 deaths under the age of 5 worldwide is due to a water-related disease [7]. Clean and safe water is essential to healthy living. Tiny worms and bacteria live in water naturally [8].

In many third-world countries, potable water is collected from communal sources, either unimproved improved; these include or unprotected wells, unprotected springs, rivers, boreholes, and public standpipes [9]. Microbial contamination of drinking water during and after collection from the water source has been recognized as a problem for households, resulting in poor health conditions and poorer water quality in storage [10,11]. According to the WHO/UNICEF report, 30% of people worldwide lack access to safe, readily drinkable water at home, and 60% lack safely managed sanitation [12]. Public health usually mentions the roles of basic hygiene and water quality that cause pathogen transmission without mentioning the gender roles that affect the overall risk of various health issues [13,14].

Guatemala is also recognized to have a crisis level of water pollution and contamination [15,16,17]. Available drinking water can be highly contaminated. People often use firewood to boil the water to purify it, leading to deforestation in the country [18]. They also inhale smoke emissions, which contribute to climate change [19].

Leptospirosis, a bacterial infection that can be spread in fresh water, is found in Guatemala. Avoid swimming in fresh, unchlorinated water, such as lakes, ponds, or rivers [20]. Without a doubt, Guatemala is currently facing a water crisis, where much of the country faces the lessthan-ideal reality of relying on contaminated water for drinking and food production. Drinking water is one of the most frequently cited concerns in research on food, energy, and water systems in Guatemala. Most research to date in Guatemala (and in international water, sanitation, and hygiene research overall) has focused on microbial risks in rural areas [21]. However, chemical water quality monitoring is often limited by lack of funding, laboratory capacity, and regulatory authority, resulting in minimal data on the occurrence and distribution of chemical contaminants in tap water [22].

Living these unhealthy environment, women in Guatemala are disproportionately affected by financial situations and living conditions compared to men [23]. This disparity is particularly pronounced in the country, where virtually entire life challenges are more severe than in industrially developed nations. Numerous factors contribute to the higher vulnerability of women in these regions [24,25]. For instance, women generally have less access to education and healthcare services, and they frequently encounter significant social and economic inequalities [26,27]. These issues are compounded by systemic barriers limiting their personal economic independence and development opportunities [28]. Consequently, women in third-world countries face complex challenges that significantly impact their overall well-being and quality of life [29].

This paper investigated typical water quality in Guatemala with documented data [30,31] and attempted to raise a realistic view of healthcare accessibility, to which women are particularly vulnerable because of their social roles and living situations, with a set of questionnaire research. Furthermore, it explores the practical lifestyle patterns of women in the country.

2. METHODOLOGY AND DATA COLLECTION

2.1 Data Collected Location

This survey data was collected from a rural area in Chimaltenango, Guatemala. Chimaltenango is a city, in southwestern Guatemala. It lies 48 km from Guatemala City, in the central highlands at an elevation of 1,790 meters above sea level. Built in 1526 at south of an old Mayan fortress, the city is popular as a market region and transportation center for its adjacent town villages. The inhabitants produce grains, sugarcane, and livestock. Chimaltenango is located on the Inter-American Highway. Its population is approximately 63,000.

2.2 Questionnaire Preparations

The water quality literature was reviewed using a methodology that employed an initial search strategy using key terms such as 'water quality,' 'Guatemala,' and 'water management.' The literature search was extended to include studies investigating the water storage method in countries to ensure a comprehensive understanding and analysis of the water situation

in Guatemala [32,33,34]. This dual focus allowed a more systemic examination of the relationship between water quality and women's health.

A pre-made questionnaire was used to assess women's living conditions. The questionnaire contained seven questions about women's general lifestyle, and thirty women participated in it. The survey questions were translated into Spanish to better understand the participants. The thirty women came from various villages to our medical volunteer sites in Guatemala to assess their medical or health issues. Asking for a survey at the medical volunteer site allowed us to simultaneously assess living conditions and their effect on women's health.

2.3 Questionnaires

Each question is given in Table 1.

Table 1. Presents the questions and answers in our questionnaire sheet.

Questions	Answers (Ranges and Choice)			
1. What is your age?"	20 - 30	32 -40	>	40
(¿Cuál es tu edad?)				
"Who do you live with?"	Parents	Hijos	Husband	Familia
(¿Con quién vives?)				
3. "At what age did you give birth to	<19	20 – 40	Ν	lo Child
your first child?				
(¿A que edad diste a luz a tu				
<u> </u>	Poor	Foir	C	Sood
health?"	FUUI	Faii	Ċ	000
(¿Cómo calificarías tu salud				
general?).				
5. "Do you face challenges	Yes		No	
accessing healthcare?"				
(¿Enfrentas desafíos para				
acceder a la atención médica?).				
"Do you face challenges	None	2 – 5 yrs	2 – 6 months	>6 months
accessing healthcare?"				
(¿Enfrentas desafíos para				
acceder a la atención médica?).				
7. "When was the last time you saw	Gynecology	Orthopedic	Gastro-	Other
a doctor, and what was the			enterology	
reason for your visit?"				
(¿Cuando fue la ultima vez que				
viste a un medico y cual fue el				
⁸ "Have you ever experienced	Voc		No	
discrimination based on your	165		INU	
dender?"				
(; Alguna vez has experimentado				
discriminación basada en tu				
género?).				

2.4 Methods of Filling the Questionnaire

A township meeting was held, while our team was there on medical service activities. The people of the town should register their name, address, and age for getting treatment from a group of health professionals. During this registration time, our survey team distributed the questionnaire paper and asked them to fill in their answers.

2.5 Data Analysis

Our survey data was carefully typed into a Microsoft Excel worksheet, and statistical procedures were performed for summary with mean and standard deviation for pie-chart plotting. The summarized answer compositions are given in percentages.

3. RESULTS

3.1 Water Quality in Guatemala

In Guatemala, there are two different types of water systems: primary drinking water and secondary water. Primary drinking water is the most frequently consumed, and secondary water is other water in households. While primary drinking water was stored in a specific 5-gallon water bottle, the secondary water was stored in the water tank for each household (Fig. 1).

Although 64% of people consume primary drinking water, 74% of people said they also drink secondary water, other than primary drinking water, once or more a week. According to the study, people who only drink primary drinking water have a significantly lower rate of GI disease than people who drink both primary drinking water and secondary water, and it was proved by the fact that secondary water has 35 % more E. colil bacteria than primary drinking water [35]. Furthermore, a study shows that about half of the studied households were drinking water containing arsenic>9 µg/L, and 13% of households exceeded the water quality standards of Guatemala and US Health [36]. Both studies showed that people in Guatemala drink water contaminated in households, which causes various GI diseases.

3.2 Surveys of Women in Guatemala

The survey conducted on women in Guatemala consists of seven questions designed to gather

comprehensive data on several aspects, including age, house composition, age at childbirth, health status, access to medical care, and awareness of rights. These questions were asked to women at various medical volunteer sites throughout Guatemala. The survey was conducted in Spanish to ensure effective communication and accurate responses.

3.2.1 Age distribution

2 presents the results of the first The Fig. question that was asked to thirty participants. The question, "What is your age?" (¿Cuál es tu edad?) was designed to collect demographic information on the age distribution of women visiting the medical tent. The data indicate that the majority of patients were in the 20 to 30 age group, comprising 46.7% of the respondents. Women aged over 40 represented the second largest group at 33.3%, while those ages 31 to 40 were the least represented at 20%. Given that some participants were mothers or grandmothers with at least two children, these findings suggest that women in Guatemala are likely to marry at a relatively young age.

3.2.2 Family member question

The Fig. 3 illustrates the findings from the second question asked to thirty participants, who inquired, "Who do you live with?" (¿Con quién vives?). This question was designed to gather information regarding the household composition of women visiting the medical tent. The data provide insight into the general household arrangement in Guatemala. The results reveal that most respondents, accounting for 33.3%, live with their entire family, including children. The second largest group, representing 30% of the respondents, consists of women who reside with their parents. Women living exclusively with their husbands constitute 23.3% of the sample, while those living solely with their children are the smallest group, at 13.3%. The results indicate that a significant number of women, precisely 43.3%, do not reside with their husbands and live with their parents or children. This data leads to the assumption that women in these households may face increased financial difficulties due to the absence of men. Consequently, based on the overall result and number of responses, the researcher also assumed that there are two predominant family structures in Guatemala: with men in the household or without men in the household.

Lim; J. Global Ecol. Environ., vol. 20, no. 3, pp. 55-67, 2024; Article no.JOGEE.12348



Fig. 1. The water tank on the ceiling of Guatemala households



Fig. 2. Shows the pie chart of age distribution (n=30).



Fig. 3. Illustrates the pie charge from the question on family members (n=30)





Fig. 4. Presents the pi-chart on the question of the age of first childbirth





3.2.3 Question on first baby born

The Fig. 4 presents the findings from the third question asked to thirty participants, which asked, "At what age did you give birth to your first child?" (¿A qué edad diste a luz a tu primer hijo?). This question was intended to gather information on the age of first childbirth among women visiting the medical tent, with the hypothesis that the age of childbirth may be linked to various health issues, such as serious infections or postpartum depression. The results indicate that nearly half of respondents, 46.7%, gave birth to their first child before the age of 19.

The second largest group, comprising 33.3% of the respondents, had their first child between 20 and 40. Additionally, approximately 20% of respondents had not yet given birth. The data offers valuable insight into the typical age of childbirth in Guatemala and aligns with the existing suggestion that women in third-world countries often marry and birth children at younger ages compared to women in developed nations. This data supports the assertion that early childbirth is prevalent among women in third-world countries, reflecting broader sociocultural patterns and health implications documented in prior research.

3.2.4 Questionnaire on overall health

The Fig. 5 presents the findings from the fourth question asked thirty participants, asking, "How would you rate your overall health?" (¿Cómo calificarías tu salud general?). This question was aimed to assess the self-awareness of women visiting the medical tent regarding their health status. The results reveal that nearly half of respondents, 46.7%, rated their health as fair, indicating it was neither good nor poor. The second largest group, comprising 33.3% of the respondents, rated their health poor. Lastly, 20% of respondents considered their overall health to be good. This finding highlights a significant awareness among the participants about their health conditions despite having limited access medical care. The data support the to assertion that women in Guatemala face substantial barriers to adequate medical treatment, likely stemming from financial challenges and limited healthcare access. This underscores the necessity for targeted interventions to improve health outcomes for women in this region.

3.3 Questionnaire on Healthcare Challenges

The Fig. 6 presents the findings from the fifth question asked by thirty participants: "Do you face challenges accessing healthcare?" (¿Enfrentas desafíos para acceder a la atención médica?). This question was designed to evaluate the practical circumstances of women who visited the medical tent regarding their

access to healthcare resources. The results indicate that 56.7% of respondents reported experiencing difficulties accessing medical care, while 43.3% did not face such challenges. These findings underscore the practical healthcare accessibility issues faced by individuals in Guatemala. The data suggest that a significant portion of the population struggles to obtain appropriate medical treatment and highlights the need for increased support from both local and national governments to improve the healthcare situation in the country.

3.4 Questionnaire on Health Professional Exposure Duration

Fig. 7 illustrates the findings from the fifth question asked by thirty participants: "Do you face challenges accessing healthcare?" (¿Enfrentas desafíos para acceder a la atención médica?). This question was designed to evaluate the practical circumstances of women who visited the medical tent regarding their access to healthcare resources. The results indicate that 56.7% of respondents reported experiencing difficulties accessing medical care, while 43.3% did not face such challenges. These findings underscore the practical healthcare accessibility issues faced by individuals in Guatemala. The data suggest that a significant portion of the population struggles to obtain appropriate medical treatment and highlights the need for increased support from both local and national governments to improve the healthcare situation in the country.



Fig. 6. Presents the pi-chart on the answer from healthcare accessibility





Fig. 7. Gives the pi-chart from the questions on doctor visiting duration





3.5 Questionnaire on the Reason of Doctor's Office Visit

The Fig. 7 present the findings from the sixth question asked by thirty participants: "When was the last time you saw a doctor, and what was the reason for your visit?" (¿Cuándo fue la última vez que viste a un médico y cuál fue el motivo de tu visita?). This question aimed to evaluate the accessibility frequency and of medical consultations and the reason for visits. By analyzing responses, the researcher assessed the regularity with which individuals sought medical attention and identified common health issues. This question provided detailed and

specific insight related to Question 5 concerning women's access to healthcare resources at the medical tent. The results indicate that 33.3% of respondents reported seeing a doctor 2 to 5 second largest years ado. The group, comprising 26.7% of respondents, indicated they had seen a doctor less than six months ago. 23.3% Approximately respondents of reported seeing a doctor between 6 months and two years ago, while 16.7% never saw doctors. These findings reveal that a significant Guatemalan population has limited access to regular medical care, underscoring the challenges faced in obtaining necessary healthcare services.



Fig. 9. Shows the pie chart on the question of gender discrimination.

То accuratelv assess women's current symptoms, the researcher inquired about the reasons for visiting the medical tent, not the reason for the previous visit to the doctor. The results show that 46.7% of respondents sought medical care for gynecological issues such as Urinary Tract Infections (UTIs) and severe menstrual pain. Additionally, 33.3% of respondents reported visiting the doctor due to Irritable Bowel Syndrome (IBS), stomach pain, or constipation. Orthopedic issues, including back, neck, and muscle pain, accounted for 6.7% of visits, while 13.3% of respondents cited other reasons for seeking medical care. The data suggest that a significant portion of the population experiences symptoms potentially linked to untreated water and dietary factors. such as UTIs, diarrhea, and stomach pain. The result highlights the need for enhanced support to improve daily living conditions and broaden access to medical care for the population in Guatemala.

3.6 Questionnaire on Gender Discrimination

The Fig. 8 presents the findings from the final question asked by thirty participants: "Have you ever experienced discrimination based on your gender?" (¿Alguna vez has experimentado discriminación basada en tu género?). This question aimed to assess the participants' self-awareness and the reality of gender discrimination in Guatemala, including issues

such as unfair wages, denial of granted rights, and various limitations due to gender. The results reveal that 63.3% of respondents reported they had not experienced gender-based discrimination, while 36.7% indicated that they had encountered such discrimination. This finding emphasizes that many women in Guatemala continue to face gender inequality and struggle to secure their entitled rights. The data support the assertion that women in Guatemala experience unfair treatment compared to women in developed nations, highlighting the need for the government and related organizations to intervene and improve conditions for women in the country.

4. DISCUSSION

The findings of this study provide information on the significant health and social challenges faced by women in third-world countries, specifically focusing on water quality, healthcare access, and gender discrimination in Guatemala. The data from our survey and literature review reveal several critical issues that require attention and intervention.

Firstly, the survey results highlight the current healthcare accessibility situation among women in Guatemala. With 56.7% of respondents indicating challenges in accessing medical care, it is evident that there are potential barriers preventing women from receiving healthcare.

These barriers may include financial limitations. geographic limitations, and insufficient healthcare institutions. The high percentage of women seeking medical attention for gynecological issues (46.7%) and gastroenterological issues (33.3%) emphasizes the untreated conditions that significantly impact women's quality of life. These findings align with previous studies documenting the adverse effects of poor water quality and inadequate sanitation on women's health in third-world countries [37]. The correlation between untreated water and health issues such as UTIs, Irritable Bowel Syndrome (IBS), and eczema, and women-specific disease, such as dysmenorrhea, suggests a direct link between environmental factors and women's health outcomes [38].

Furthermore, the results regarding gender discrimination reveal that 36.7% of women have experienced gender-based discrimination. This statistic is alarming, considering the various forms of discrimination reported, including unfair wages, denial of rights, and suppression of voices. These forms of discrimination not only violate fundamental human rights but also contribute to the situation of gender inequality, limiting women's opportunities for education, employment, and overall well-being.

5. CONCLUSION

The research provides critical insights into the various issues women face in Guatemala. particularly regarding water guality and health issues. The findings highlight that poor water quality significantly affects women's health, leading to conditions such as urinary tract infections, irritable bowel syndrome, and skin issues. These health challenges are exacerbated by the socio-economic conditions in these regions, including limited access to education, healthcare, and financial opportunities. The study also highlights the urgent need for intervention to improve healthcare access, water quality, and gender equality in Guatemala. The findings advocate for comprehensive policies that address both the immediate health needs of women and the broader social determinants of health. Future research should focus on studies to monitor the impact of these policies over time and further explore the interdisciplinarity of health and environmental factors.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

The author hereby declares that NO generative Al technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

COMPETING INTERESTS

Author has declared that no competing interests exist.

REFERENCES

- 1. Bidaisee S. The importance of clean water. Biomedical Journal of Scientific & Technical Research. 2018;8. DOI:10.26717/BJSTR.2018.08.001719.
- Khalifa M, Bidaisee S. The importance of clean water. Sch J Appl Sci Res. 2018;1(7):17-20.
- Hutton G, Chase C. Water Supply, Sanitation, and Hygiene. In: Mock CN, Nugent R, Kobusingye O, et al. editors. Injury prevention and environmental health. 3rd edition. Washington (DC): The International Bank for Reconstruction and Development / The World Bank Chapter 9; 2017.

Available:https://www.ncbi.nlm.nih.gov/boo ks/NBK525207/

DOI: 10.1596/978-1-4648-0522-6_ch9.

- 4. United Nations. Conference on the midterm comprehensive review of the implementation of the objectives of the international decade for action, "Water for Sustainable Development", 2018–2028 New York; 2023.
- Sarah D, Sara G. Inequalities in water, sanitation and hygiene: Challenges and opportunities for measurement and monitoring, Water Security. 2023;20: 100143. ISSN 2468-3124, Available:https://doi.org/10.1016/j.wasec.2 023.100143.
- Shayo GM, Elimbinzi E, Shao G, Fabian C. Severity of waterborne diseases in developing countries and the effectiveness of ceramic filters for improving water quality. Bulletin of the National Research Centre; 2023;47. DOI:10.1186/s42269-023-01088-9.

- 7. UNICEF. Triple threat how disease, climate risks, and unsafe water, sanitation and hygiene create a deadly combination for children. New York; United Nations Children's Fund (UNICEF); 2023.
- Cabral JP. Water microbiology. Bacterial pathogens and water. Int J Environ Res Public Health. 2010;7(10): 3657-703. DOI: 10.3390/ijerph7103657. Epub 2010

OCI: 10.3390/ijerph/103657. Epub 2010 Oct 15. PMID: 21139855; PMCID: PMC2996186.

 WHO. Safely managed drinking water thematic report on drinking water 2017. Geneva, Switzerland: World Health Organization; 2017. CIP data are available at

http://apps.who.int/iris

10. WHO, (). Guidelines for drinking-water quality: Fourth edition incorporating the first and second addenda. Geneva: World Health Organization; 7, Microbial aspects; 2022.

Available:https://www.ncbi.nlm.nih.gov/boo ks/NBK579466/

11. Rameck M, Margaret M, Tendai K, Demetria FG. The effect of household practices on the deterioration of microbial quality of drinking water between source and point of use in Murewa district, Zimbabwe Journal of Water and Health. 2022;20(3):518.

DOI: 10.2166/wh.2022.251

- 12. UNICEF. New report on inequalities in access to water, sanitation and hygiene also reveals more than half of the world does not have access to safe sanitation services, New York, Geneva; 2019
- Jalali R. The Role of Water, Sanitation, Hygiene, and Gender Norms on Women's Health: A Conceptual Framework. Gendered Perspectives on International Development. 2021;1:21-44. Available:https://dx.doi.org/10.1353/gpi.20 21.0001.
- Pieters MM, Fahsen N, Craig C, Quezada R, Pratt CQ, Gomez A, Brown TW, Kossik A, McDavid K, Vega Ocasio D, Lozier MJ, Cordón-Rosales C. Assessment of water, sanitation, and hygiene conditions in public elementary schools in Quetzaltenango, Guatemala, in the Context of the COVID-19 Pandemic. Int J Environ Res Public Health. 2023;20(20):6914.

DOI:10.3390/ijerph20206914.

PMID: 37887652; PMCID: PMC10606716.

- 15. World Bank. Guatemala's water supply, sanitation, and hygiene poverty diagnostic: Challenges and Opportunities. WASH Poverty Diagnostic. World Bank, Washington, DC; 2018.
- Illmer P. Water and Socio-Environmental Crisis in Guatemala City's Metropolitan Area. Latin American Perspectives. 2022; 49(6):55-70. Available:https://doi.org/10.1177/0094582X 221120014
- Kondash AJ, Herrera I, Castellanos E, Baker J, Leiva B, Houtven GV, Fuentes G, Alfaro G, Henry C, Wade C, Redmon H. Food, energy, and water nexus research in Guatemala – A systematic literature review. Environmental Science & Policy. 2021;124:175-185 ISSN 1462-9011. Available:https://doi.org/10.1016/j.envsci.2 021.06.009.
- Bevan AC. A worldwide water revolution starts in Guatemala. New Internationalist, The World Unspun; 2013. Available:https://newint.org/blog/2013/08/2 0/ceramic-pot-filtration.
- Afifa Kashaf A., Nazim H, Muhammad HA, Muhammad ZS. Air pollution and climate change as grnd challenges to sustainability,Science of The Total Environment. 2024;928:172370. ISSN 0048-9697.

Available:https://doi.org/10.1016/j.scitotenv .2024.172370.

- Wynwood SJ, Graham GC, Weier SL, Collet TA, McKay DB, Craig SB. Leptospirosis from water sources. Pathog Glob Health. 2014 Oct;108(7):334-8.
 DOI:10.1179/2047773214Y.0000000156.
 Epub 2014 Oct 27. PMID: 25348115; PMCID: PMC4241785.
- Hoponick RJ, Mulhern RE, Castellanos E, Wood E, McWilliams A, Herrera I, Liyanapatirana C, et al. Participatory Science Approach to Evaluating Factors Associated with the Occurrence of Metals and PFAS in Guatemala City Tap Water. Int J Environ Res Public Health. 2022 May 15;19(10):6004.

DOI: 10.3390/ijerph19

106004. PMID: 35627541; PMCID: PMC9142002.

22. Zulkifli SN, Rahim HA, Lau WJ. Detection of contaminants in water supply: A review

on state-of-the-art monitoring technologies and their applications. Sens Actuators B Chem. 2018 Feb;255:2657-2689.

DOI: 10.1016/j.snb.2017.09.078. Epub 2017 Sep 18. PMID: 32288249; PMCID: PMC7126548.

- 23. IMF. Gender Inequality and Economic Growth: Evidence from Industry-Level Data1 Prepared by Ata Can Bertay, Ljubica Dordevic, and Can Sever Authorized for distribution by Nigel Jenkinson, JEL Classification Numbers: O40, J16, O1, O47; 2020.
- 24. UN. Women's rights are human rights: United Nations Publication sales no. E.14.XIV.5 ISBN 978-92-1-154206-6 E-ISBN 978-92-1-056789-3, New York and Genova; 2023.
- 25. Kruk ME, Gage AD, Arsenault C, Jordan K, Leslie HH, et. al. High-quality health systems in the Sustainable Development Goals era: time for a revolution. Lancet Glob Health. 2018;6(11): e1196-e1252.

DOI: 10.1016/S2214-109X(18)30386-3.

 Azad AD, Charles AG, Ding Q, Trickey AW, Wren SM. The gender gap and healthcare: associations between gender roles and factors affecting healthcare access in Central Malawi, June-August 2017. Arch Public Health. 2020;78(1): 119.

DOI: 10.1186/s13690-020-00497-w.

- Lwamba E, Shisler S, Ridlehoover W, Kupfer M, Tshabalala N, Nduku P, Langer L, Grant S, Sonnenfeld A, Anda D, Eyers J, Snilstveit B. Strengthening women's empowerment and gender equality in fragile contexts towards peaceful and inclusive societies: A systematic review and meta-analysis. Campbell Syst Rev. 2022;18(1):e1214.
 DOI: 10.1002/cl2.1214. PMID: 36913184; PMCID: PMC8904729.
- Chanda T, Ngulube L. Women in Leadership: Examining Barriers to Women's Advancement in Leadership Positions. Asian Journal of Advanced Research and Reports. 2024;18:273-290. DOI:10.9734/ajarr/2024/v18i6671.
- 29. UN Women. Gender equality and sustainable development, 2014, United Nations, The Research and Data Section of UN Women; 2014.

 Murphy Daniel E, et al. An assessment of household water quality among Peace Corps volunteers in Guatemala. Tropical Diseases, Travel Medicine and Vaccines; 2019.

Available:https://doi.org/10.1186/s40794-019-0078-8.

- Marcillo CE, et al. Drinking water quality and consumer perceptions at the point-ofuse in San Rafael Las Flores, Guatemala. Water Practice and Technology. 2020;15(2):374-385, Available:https://doi.org/10.2166/wpt.2020. 025.
- Redmon JH, Mulhern RE, Kondash AA, Lopez EJC. Understanding water quality at the tap in Guatemala City, RTI International; August 05, 2022. Available: https://www.rti.org/insights/understandingwater-quality-tap-guatemala-city
- Johnke K. Navigating water quality: Illuminations from Solola, Guatamala, Center for Reginal Food Studies, College of Social & Behavioral Sciences, The University of Arizona; 2024. Available:https://crfs.arizona.edu/news/nav igating-water-quality-illuminations-sololaguatemala.
- Interactive Country Fiches. Discovery a country through its environmental situation data. UN Environment Programme of Work Sub-programme 7 Environment under review; 2024.

Available: https://dicf.unepgrid.ch/

35. World Health Organization. WHO Global water, sanitation and hygiene annual report 2019. World Health Organization (WHO); 2020.

Available:https://www.who.int/publications/i /item/9789240013391.

 Silbergeld EK, Jodi AF. Environmental Exposures and Women's Health. Clinical Obstetrics and Gynecology. 2002;4(4): 1119-1128.
Available:https://journals.lww.com/clinicalo

bgyn/citation/2002/12000/environmental_e xposures_and_women_s_health.19.aspx.

 Kayser GL, Rao N, Jose R, Raj A. Water, sanitation and hygiene: measuring gender equality and empowerment. Bull World Health Organ. 2019;97(6):438-440. Lim; J. Global Ecol. Environ., vol. 20, no. 3, pp. 55-67, 2024; Article no.JOGEE.12348

DOI: 10.2471/BLT.18.223305. Epub 2019 May 14. PMID: 31210683; PMCID: PMC6560376.

38. Salari-Moghaddam A, Hassanzadeh KA, Esmaillzadeh A, Adibi P. Water

consumption and prevalence of irritable bowel syndrome among adults. PLoS One. 2020;15(1):e0228205.

DOI:10.1371/journal.pone.0228205. PMID: 31978193; PMCID: PMC6980581.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history: The peer review history for this paper can be accessed here: https://prh.ikprress.org/review-history/12348